


Safer CRISPR gene editing with fewer off-target hits

July 9 2020



Mutating the enzyme at the heart of the CRISPR gene editing system can improve its fidelity. Credit: DataBase Center for Life Science (DBCLS)

The CRISPR system is a powerful tool for the targeted editing of genomes, with significant therapeutic potential, but runs the risk of inappropriately editing "off-target" sites. However, a new study

publishing July 9, 2020 in the open-access journal *PLOS Biology* by Feng Gu of Wenzhou Medical University, China, and colleagues, shows that mutating the enzyme at the heart of the CRISPR gene editing system can improve its fidelity. The results may provide a therapeutically safer strategy for gene editing than using the unmodified enzyme system.

The CRISPR system employs an [enzyme](#) called Cas9 to cleave DNA. Cas9 will cut almost any DNA sequence. Its specificity comes from its interaction with a "guide RNA" (gRNA) whose sequence allows it to bind with the target DNA through base-pair matching. Once it does, the enzyme is activated and the DNA is cut.

The CRISPR system is found in multiple [bacterial species](#); among those commonly used in research, that from *Staphylococcus aureus* has the advantage of size—unlike some others, its gene is small enough to fit inside a versatile and harmless gene therapy vector called adeno-associated virus, making it attractive for therapeutic purposes.

A key limitation of any of the CRISPR systems, including that from *S. aureus*, is off-target cleavage of DNA. A guide RNA may bind weakly to a site whose sequence is a close but imperfect match; depending on how close the match is and how tightly the enzyme interacts with the paired gRNA-DNA complex, the enzyme may become activated and cut the DNA wrongly, with potentially harmful consequences.

To explore whether the *S. aureus* Cas9 could be modified to cleave with higher fidelity to the intended target, the authors generated a range of novel Cas9 mutants and tested their ability to discriminate against imperfect matches while retaining high activity at the intended site. They found one such mutant, which distinguished and rejected single base-pair mismatches between gRNA and DNA, regardless of the target, increasing the fidelity up to 93-fold over the original enzyme. They showed that the mutation affected part of the recognition domain, the

region of the enzyme that coordinates contacts between the enzyme and the gRNA-DNA complex. The mutation had the likely effect of weakening those contacts, thus ensuring that only the strongest pairing—which would come from a perfect sequence match—would trigger enzyme activity.

"Avoidance of off-target cleavage is a crucial challenge for development of CRISPR for medical interventions, such as correcting genetic diseases or targeting cancer cells," Gu said. "Our results point the way to developing potentially safer gene therapy strategies."

More information: Xie H, Ge X, Yang F, Wang B, Li S, Duan J, et al. (2020) High-fidelity SaCas9 identified by directional screening in human cells. *PLoS Biol* 18(7): e3000747.
doi.org/10.1371/journal.pbio.3000747

Provided by Public Library of Science

Citation: Safer CRISPR gene editing with fewer off-target hits (2020, July 9) retrieved 9 May 2024 from <https://phys.org/news/2020-07-safer-crispr-gene-off-target.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.