


Team simultaneously synthesizes dicarboxylic acids and hydrogen from diols

July 30 2020



Using an iridium catalyst bound to a bipyridonate ligand, researchers were able to synthesize dicarboxylic acids from aqueous diols, with the added benefit of generating hydrogen as a byproduct. Credit: Kyoto University

Every year, the chemical industry makes trillions of dollars synthesizing the countless chemical compounds we use every day.

From pharmaceuticals for keeping you healthy to polyester woven into your shirt, [industrial processes](#) turn simple chemicals into complex,

valuable [compounds](#). Researchers in turn work constantly to develop new substances as well as safer and more efficient methods of production.

In a paper published in *ChemSusChem*, researchers from Kyoto University's Graduate School of Human and Environmental Studies describe how they have significantly improved the process that leads to polymers and plasticizers. And as an added bonus, their method generates hydrogen as a byproduct.

"We use renewable [chemical](#) compounds collectively known as *diols* to dramatically change the synthesis of [dicarboxylic acids](#)," explains corresponding author Ken-ichi Fujita, "These are needed to produce polymers, plasticizers, and lubricants."

Existing production methods, based on the oxidation of hydrocarbons using toxic oxidants, generate harmful waste products. Knowledge of this motivated the team to work on a new method of synthesis, changing the initial compound along with the catalyst.

"We began by looking at liquid solutions of *diols*, which are more renewable than starting materials previously used," continues Fujita.

"Then we needed to find a catalyst, and we settled on using iridium bound to a bipyridonate ligand."

The team were pleased to find that combining the *diols* with the new catalyst generated dicarboxylic acids with greater efficiency and significantly more yield, as well as four equivalents of hydrogen for every unit of *diols*.

Developing more efficient and safer methods of compound production is vital for industrial organic chemistry, and for the global environment.

"It is a pleasant surprise to see how efficient our new process is," concludes Fujita. "We hope to continue to improve safety and efficiency as we discover alternative chemical processes that are valuable to all humanity."

More information: Genki Toyooka et al, Synthesis of Dicarboxylic Acids from Aqueous Solutions of Diols with Hydrogen Evolution Catalyzed by an Iridium Complex, *ChemSusChem* (2020). [DOI: 10.1002/cssc.202001052](https://doi.org/10.1002/cssc.202001052)

Provided by Kyoto University

Citation: Team simultaneously synthesizes dicarboxylic acids and hydrogen from diols (2020, July 30) retrieved 20 April 2024 from <https://phys.org/news/2020-07-team-simultaneously-dicarboxylic-acids-hydrogen.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.