

Bacteria can defuse dangerous chemical in Passaic River

August 19 2020, by Todd Bates



A view of the dioxin-contaminated Passaic River in Newark, New Jersey. Credit: Donna E. Fennell

Bacteria that can help defuse highly toxic dioxin in sediments in the Passaic River—a Superfund hazardous waste site—could eventually aid cleanup efforts at other dioxin-contaminated sites around the world, according to Rutgers scientists.

Their research, published in the journal *Environmental Science & Technology*, needs further work to realize the full potential of the beneficial bottom-dwelling microbes.

"The bacteria-driven process we observed greatly decreases the toxicity of [dioxin](#)," said senior author Donna E. Fennell, a professor who chairs the Department of Environmental Sciences in the School of Environmental and Biological Sciences at Rutgers University–New Brunswick.

"Our results showed that although the process is quite slow, it can be enhanced and may even have the potential to remove all toxic chlorines from the compound," said lead author Rachel K. Dean, a Rutgers doctoral student.

In a process known as dechlorination, the bacteria remove chlorine atoms from 2,3,7,8-Tetrachlorodibenzo-p-dioxin (TeCDD), the most toxic of the many dioxins, according to the study.

These chemicals can cause cancer, reproductive and developmental problems and immune system damage, according to the U.S. Environmental Protection Agency. They also can interfere with hormones in the body.

In New Jersey, the decades-old Diamond Alkali Superfund site includes a former chemical manufacturing facility in Newark, a 17-mile tidal stretch of the Passaic River and tributaries, Newark Bay and portions of the Hackensack River, Arthur Kill and Kill van Kull, according to the

EPA.

The 2,3,7,8-TeCDD dioxin is a byproduct of combustion and chemical product manufacturing, including the herbicides in Agent Orange. Sampling revealed high levels of dioxin in 1983 and the site landed on the Superfund National Priorities List a year later. Dioxin, polychlorinated biphenyls, metals, [polycyclic aromatic hydrocarbons](#) and pesticides were found in sediment in the Lower Passaic River.

Though dredging is required to remove the most highly contaminated sediments in the Passaic River, some contamination has spread and will remain in the river and estuary, where it could be transformed by the bacteria over time, according to the Rutgers scientists.

In the study, scientists took solid material from the river bottom and mixed it with water and other nutrients in the lab to make mud in bottles. Then they added 2,3,7,8-TeCDD and another chemical (dichlorobenzene) that boosted dechlorination by bacteria.

The Rutgers scientists revealed bacteria that are likely involved in the dechlorination process—a novel Dehalococcoidia bacterium—by sequencing their DNA from the dioxin-tainted mud. It is possible that 2,3,7,8-TeCDD could be completely dechlorinated in the Passaic River and estuary.

The next goal is to try to identify the enzymes involved in dechlorination so cleanup technologies could be developed that lead to more dechlorination at this and other contaminated sites, the study says.

More information: Rachel K. Dean et al, 2,3,7,8-Tetrachlorodibenzo-p-dioxin Dechlorination is Differentially Enhanced by Dichlorobenzene Amendment in Passaic River, NJ Sediments, *Environmental Science & Technology* (2020). [DOI: 10.1021/acs.est.0c00876](https://doi.org/10.1021/acs.est.0c00876)

Provided by Rutgers University

Citation: Bacteria can defuse dangerous chemical in Passaic River (2020, August 19) retrieved 5 May 2024 from <https://phys.org/news/2020-08-bacteria-defuse-dangerous-chemical-rassaic.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.