

## Why black rhinos may get sick in captivity

August 12 2020



Credit: Pixabay/CC0 Public Domain

Inflammation and oxidative stress may be involved in the pathogenesis of iron overload disorder in captive black rhinoceroses, making this syndrome a potential common denominator to various diseases described in captivity in this species, according to a study published August 12 in the open-access journal *PLOS ONE* by Hanae Pouillevet of Oniris



Nantes-Atlantic National College of Veterinary Medicine, and colleagues.

Black rhinoceroses are considered to be vulnerable to critically endangered by the International Union for Conservation of Nature. In captivity, these animals develop serious diseases not seen in wild black rhinoceroses. One such condition is iron overload disorder, in which iron accumulates in the body, potentially contributing to organ failure. In the new study, the researchers tested how iron overload disorder relates to inflammation and oxidative stress, which have been linked to a variety of health conditions. Inflammation is part of the complex biological response of body tissues to protect against harmful stimuli, such as pathogens, damaged cells, or irritants. Oxidative stress is a phenomenon caused by an imbalance between the production of reactive oxygen species in cells and tissues and the ability to detoxify these oxygencontaining, chemically reactive molecules through protective mechanisms.

15 black rhinoceroses and 29 white rhinoceroses were sampled at 22 European zoos. Compared to white rhinoceroses, which do not develop iron overload disorder, black rhinoceroses showed higher iron levels and higher markers of inflammation and oxidative stress. The findings suggest that iron accumulation may enhance inflammation and oxidative stress, leading to additional diseases, rapid aging, and organ damage. According to the authors, the results highlight the importance of continuing efforts to control iron overload disorder in <u>captive</u> black rhinoceroses, whether through dietary interventions or through regular blood removal. Further investigations are needed to assess the value of the inflammatory and oxidative markers for determining the prognosis of captive black rhinoceroses, particularly for evaluating the impact of reduced-<u>iron</u> and antioxidant-supplemented diets.

More information: Pouillevet H, Soetart N, Boucher D, Wedlarski R,



Jaillardon L (2020) Inflammatory and oxidative status in European captive black rhinoceroses: A link with Iron Overload Disorder? *PLoS ONE* 15(8): e0231514. doi.org/10.1371/journal.pone.0231514

## Provided by Public Library of Science

Citation: Why black rhinos may get sick in captivity (2020, August 12) retrieved 23 June 2024 from <u>https://phys.org/news/2020-08-black-rhinos-sick-captivity.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.