


# Hot sulfur breath: Extremophilic archaea provide clues on evolution of sulfur metabolism

August 24 2020

---



Credit: CC0 Public Domain

The use of sulfur compounds to produce energy is one of the most ancient types of metabolisms used by primitive microorganisms to thrive on the early anoxic Earth. How this microbial lifestyle has been evolving remains unclear.

The increase of oxygen in the atmosphere approximately 2.5 billion years ago lead to the accumulation of [sulfate](#), the most-oxidized form of sulfur. Sulfate reduction became the most important process in the global cycle of sulfur. It is a dominant microbial process in oxygen-free environments, such as marine and freshwater sediments. However, the geological evidence of microbial dissimilatory sulfate reduction (DSR) dates back to about 3.5 billion years ago, and predates the appearance of oxygen in the atmosphere.

Since the [early life](#) was evolving under hot and anoxic conditions, it was therefore important to examine the microorganisms from modern ecosystems with similar conditions in the environments resembling those on early Earth. The study was therefore focussed on a new archaeon "Candidatus *Vulcanisaeta moutnovskia*," an acid- and hot-temperatures-loving microorganism from acidic hot spring in Kamchatka Peninsula (Russia).

In an article published in *Nature Microbiology*, an international team of researchers led by Dr. Nikolay Chernyh and Prof Elizaveta Bonch-Osmolovskaya of the Winogradsky Institute of Microbiology in Moscow, colleagues from Madrid, Vienna and Bangor University's Center for Environmental Biotechnology (CEB) at the School of Natural Sciences revealed that the sulfate reduction in the hot spring is attributed to "Candidatus *Vulcanisaeta moutnovskia*." The team used microbial community profiling coupled with growth experiments, radioisotopes and proteomics and analysis of evolutionary trajectory of DSR genes.

Sulfate reduction in Archaea is less known, as compared with Bacteria, the most numerous group of living organisms. However, the analysis of genomes suggests that the genes for the second reaction in DSR are more ancient in Archaea, and were probably even present in the hypothetical "Last Universal Common Ancestor" (LUCA) organism.

"Evidently, the sulfate reduction in the hot spring is facilitated by 'Vulcanisaeta' that have recruited some genetic loci for this process from bacteria, their neighbors in the environment, a very long time ago. 'Vulcanisaeta' can possibly be considered a relic form of life in such environments and is a great model organism to study the evolution of sulfur metabolism," comments Prof Peter Golyshin from CEB, Bangor University, who contributed to the analysis of proteomic data in this study.

**More information:** Nikolay A. Chernyh et al. Dissimilatory sulfate reduction in the archaeon 'Candidatus Vulcanisaeta moutnovskia' sheds light on the evolution of sulfur metabolism, *Nature Microbiology* (2020). DOI: [10.1038/s41564-020-0776-z](https://doi.org/10.1038/s41564-020-0776-z)

Provided by Bangor University

Citation: Hot sulfur breath: Extremophilic archaea provide clues on evolution of sulfur metabolism (2020, August 24) retrieved 8 September 2024 from <https://phys.org/news/2020-08-hot-sulfur-extremophilic-archaea-clues.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.