

Magnetized gas flows feed a young star cluster

August 18 2020





Composite image of the Serpens South Cluster. Magnetic fields observed by SOFIA are shown as streamlines over an image from the Spitzer Space Telescope. SOFIA indicate that gravity can overcome some of the strong magnetic fields to deliver material needed for new stars. The magnetic fields have been dragged into alignment with the most powerful flows, as seen in the lower left where the streamlines are following the direction of the narrow, dark filament. This is accelerating the flow of material from interstellar space into the cloud, and fueling the collapse needed to spark star formation. Credit: NASA/SOFIA/T. Pillai/J. Kauffmann; NASA/JPL-Caltech/L. Allen

Observations of magnetic fields in interstellar clouds made of gas and dust indicate that these clouds are strongly magnetized, and that magnetic fields influence the formation of stars within them. A key observation is that the orientation of their internal structure is closely related to that of the magnetic field.

To understand the role of magnetic fields, an international research team led by Thushara Pillai, Boston University & Max Planck Institute for Radio Astronomy (MPIfR) in Bonn, Germany, observed the filamentary network of the dense gas surrounding a young star cluster in the solar neighboorhood, with the HAWC+ polarimeter on the airborne observatory SOFIA at infrared wavelengths. Their research shows that not all dense filaments are created equal. In some of the filaments the <u>magnetic field</u> succumbs to the flow of matter and is pulled into alignment with the filament. Gravitational force takes over in the denser parts of some filaments and the resulting weakly magnetized gas flow can feed the growth of young stellar clusters like a conveyor belt.

The results are published in this week's issue of Nature Astronomy.

The interstellar medium is composed of tenuous gas and dust that fills the vast amount of emptiness between stars. Stretching across the



Galaxy, this rather diffuse material happens to be a significant mass reservoir in Galaxies. An important component of this interstellar gas are the cold and dense molecular clouds which hold most of their mass in the form of molecular hydrogen. A major finding in the last decade has been that extensive network of filaments permeate every molecular cloud. A picture has emerged that stars like our own sun form preferentially in dense clusters at the intersection of filaments.

The researchers observed the filamentary network of dense gas around the Serpens South Cluster with HAWC+, a polarization-sensitive detector onboard the airborne observatory SOFIA, in order to understand the role of magnetic fields. Located about 1,400 light-years away from us, the Serpens South cluster is the youngest known cluster in the local neighborhood at the center of a network of dense filament.

The observations show that low-density gaseous filaments are parallel to the magnetic field orientation, and that their alignment becomes perpendicular at higher gas densities. The high angular resolution of HAWC+ reveals a further, previously unseen twist to the story. "In some dense filaments the magnetic field succumbs to the flow of matter and and is pulled into alignment with the filament," says Thushara Pillai (Boston University and MPIfR Bonn), the first author of the publication. "Gravitational force takes over in the more opaque parts of certain filaments in the Serpens Star Cluster and the resulting weakly magnetized gas flow can feed the growth of young stellar clusters like a conveyor belt," she adds.

It is understood from theoretical simulations and observations that the filamentary nature of molecular clouds actually plays a major role in channeling mass from the larger interstellar medium into young stellar clusters whose growth is fed from the gas. The formation and evolution process of stars is expected to be driven by a complex interplay of several fundamental forces—namely turbulence, gravity, and the



magnetic field. In order to get an accurate description for how dense clusters of stars form, astronomers need to pin down the relative role of these three forces. Turbulent gas motions as well as the mass content of filaments (and therefore gravitation force) can be gauged with relative ease. However, the signature of the interstellar magnetic field is weak, also because it is about 10,000–times weaker than even our own Earth's magnetic field. This has made measurements of magnetic field strengths in filaments a formidable task.

"The magnetic field directions in this new polarization map of Serpens South align well with the direction of gas flow along the narrow southern filament. Together these observations support the idea that filamentary accretion flows can help form a young star <u>cluster</u>," adds Phil Myers from the Harvard-Smithsonian Center for Astrophysics, a co-author of the paper.

A small fraction of a molecular cloud's mass is made up by small dust grains that are mixed into the interstellar gas. These interstellar dust grains tend to align perpendicular to the direction of the magnetic field. As a result, the light emitted by the dust grains is polarized—and this polarization can be used to chart the magnetic field directions in molecular clouds.

Recently, the Planck space mission produced a highly sensitive all–sky map of the polarized dust emission at wavelengths smaller than 1 mm. This provided the first large–scale view of the magnetization in filamentary molecular clouds and their environments. Studies done with Planck data found that filaments are not only highly magnetized, but they are coupled to the magnetic field in a predictable way. The orientation of the magnetic fields is parallel to the filaments in low–density environments. The magnetic fields change their orientation to being perpendicular to filaments at high gas densities, implying that magnetic fields play an important role relative in shaping filaments,



compared to the influence of turbulence and gravity.

This observation pointed towards a problem. In order to form stars in gaseous filaments, the filaments have to lose the magnetic fields. When and where does this happen? With the order of magnitude higher angular resolution of the HAWC+ instrument in comparison to Planck it was now possible to resolve the regions in filaments where the magnetic filament becomes less important.

"Planck has revealed new aspects of magnetic fields in the <u>interstellar</u> <u>medium</u>, but the finer angular resolutions of SOFIA's HAWC+ receiver and ground-based NIR polarimetry give us powerful new tools for revealing the vital details of the processes involved," says Dan Clemens, Professor and Chair of the Boston University Astronomy Department, another co-author.

"The fact that we were able to capture a critical transition in star formation was somewhat unexpected. This just shows how little is known about cosmic magnetic fields and how much exciting science awaits us from SOFIA with the HAWC+ receiver," concludes Thushara Pillai.

More information: Thushara G.S. Pillai et al. Magnetized filamentary gas flows feeding the young embedded cluster in Serpens South, *Nature Astronomy* (2020). DOI: 10.1038/s41550-020-1172-6

Provided by Max Planck Society

Citation: Magnetized gas flows feed a young star cluster (2020, August 18) retrieved 17 May 2024 from <u>https://phys.org/news/2020-08-magnetized-gas-young-star-cluster.html</u>



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.