

# Out of sync: Ecologists report climate change affecting bee, plant life cycles

August 19 2020, by Mary-Ann Muffoletto

---



*Andrena transnigra* (black-banded miner bee) on *Solidago multiradiata* (Rocky Mountain goldenrod) at the Rocky Mountain Biological Laboratory near Gothic, Colorado. Utah State University and North Carolina State University ecologists, along with a multi-institution team, report on a community-wide assessment revealing phenological mismatch between bees and flowering plants, due to

climate change, in the August 19, 2020 issue of *'Ecology Letters.'* Credit: David Inouye, RMBL

Bees and flowers seem inseparable harbingers of spring, but what happens when pollinators emerge later than their sources of nectar and pollen? Reporting on the first community-wide assessment of 67 bee species of the Colorado Rockies, ecologists Michael Stemkovski of Utah State University and Rebecca Irwin of North Carolina State University say "phenological mismatch," changing timing of life cycles between bees and flowers, caused by climate change, has the potential to disrupt a mutually beneficial relationship.

"We analyzed time-series abundance data collected at 18 sites around the Rocky Mountain Biological Laboratory (RMBL) in the Elk Mountains of western Colorado during a nine-year, National Science Foundation-funded bee monitoring project," says Stemkovski, doctoral student in USU's Department of Biology and the USU Ecology Center.

He and Irwin, senior author, along with colleagues from RMBL, University of Texas at Austin, Imperial College London, University of Manitoba, USDA-ARS Pollinating Insects Research Unit at USU, Central Texas Melittological Institute, Royal Saskatchewan Museum, Texas A&M University, Florida State University and University of Maryland, published findings in the August 19, 2020 issue of *Ecology Letters*.

"We find bee emergence timing is advancing with snowmelt timing, but bee phenology—timing of emergence, peak abundance and senescence—is less sensitive than flower phenology," says Irwin, professor of applied ecology at NCSU. "Given global concerns about pollinator declines, the research provides important insight into the

potential for reduced synchrony between flowers and their pollinators under climate change."

Previous studies focused primarily on temperature, Stemkovski says, but this study probed the effects of topography and bee species traits, as well.

"Elevation played a large role in when bees start foraging, as well as the bees' functional traits, such as whether bees nested below or above ground, and the life stage in which they overwintered," he says. "We found all of these factors predicted bee emergence, but the most important factor was snowmelt timing."

If bees begin foraging later than spring plants reach their flowering peak, consequences could be reduced abundance of pollinators, from limited sustenance, and reduced abundance of plants, from limited pollination.

"In the short-term, we expect mutualist species to suffer fitness losses," Stemkovski says. "In the long-term, [bees](#) and plants may be able to adapt and reestablish some synchrony, unless [climate change](#) outpaces the rate of adaptation."

**More information:** "Bee phenology is predicted by climatic variation and functional traits," *Ecology Letters* (2020). [DOI: 10.1111/ele.13538](https://doi.org/10.1111/ele.13538)

Provided by Utah State University

Citation: Out of sync: Ecologists report climate change affecting bee, plant life cycles (2020, August 19) retrieved 25 May 2024 from <https://phys.org/news/2020-08-sync-ecologists-climate-affecting-bee.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.