


Velcro method for more precise binding of drug particles

August 24 2020



TU/e researchers prove selectivity based on number of receptors on the cell

walls. Credit: TU Eindhoven

In order to deliver drug particles to the right place in the body—a field known as nanomedicine—selectivity plays an important role. After all, the drug only has to attach itself to the cells that need it. A theory from 2011 predicts that selectivity is not only based on the type of receptor, but also on the number and strength of the receptors on the cell. Researchers at Eindhoven University of Technology are now proving this experimentally. They have published their results in the journal *PNAS*.

Cells interact with each other through [receptors](#) and ligands. They fit on each other like a key in a lock; a ligand of one cell only fits on the appropriate target receptor of the other cell. The field of nanomedicine makes use of this by imitating ligands that fit the receptors of the diseased cell that needs the drug.

In 2011, Daan Frenkel and his group in Cambridge used a theoretical model to predict that not only the type of ligands and receptors play a major role, but also the number and strength. This means that even weak ligands can bind, as long as there are enough receptors present on the surface of the target cell. Researchers Max Scheepers, Leo van IJzendoorn, and Menno Prins, all part of the Institute for Complex Molecular Systems, have now proven this theory experimentally with particles for the first time.

Many weak bonds become strong

Van IJzendoorn: "Compare it to Velcro. If one hook is fastened, the strip does not stick immediately. Only when several hooks are fastened does the bond become strong enough. This is also how it works in the human

body; the weak binding of a ligand on a receptor becomes enormously strong the more there are."

And that's a useful feature for nanomedicine. Diseased [cells](#) do not always have different receptors than healthy cells, but they often have more receptors on their cell walls. By developing the drug in such a way that it only sticks to cells with a lot of receptors, you can still distinguish between diseased and healthy cells. This makes it possible to send the drug particles more precisely to the diseased cells in the body.

Single strand DNA as receptor and ligand

"We have now experimentally demonstrated with particles that many weak ligands give a high selectivity: The particles only bind if there are exactly enough receptors present. This creates a threshold value," explains van IJzendoorn. The researchers carried out a binding experiment for this purpose, designing particles with either receptor DNA or ligand DNA on its surface.

A [magnetic field](#) first pulled the particles toward each other, and after some time, released them. Van IJzendoorn: "This allowed us to optically measure how many particles had developed a strong molecular binding with each other."

By varying the number of DNA molecules and the strength of the [ligand](#)-receptor binding, not only were the researchers able to see how many bindings were needed for the particles to stay bonded, but also to observe the emergence of the [threshold value](#).

Nanomedicine and biosensors

Van IJzendoorn says, "These results form a new benchmark for

understanding and applying selectivity in biomedical applications. The work provides a fundamental basis for the design of binding processes in nanomedicine. In addition, it is important for the development of nanotechnological biosensors, because particles are also used in these systems for the establishment of selective bonds."

This research was published on 24 August in the journal *PNAS*, titled "Multivalent weak interactions enhance selectivity of inter-particle binding." The research was carried out at Eindhoven University of Technology, at the departments of Applied Physics and Biomedical Engineering and the Institute for Complex Molecular Systems.

More information: M. R. W. Scheepers et al., Multivalent weak interactions enhance selectivity of interparticle binding, *PNAS* (2020). www.pnas.org/cgi/doi/10.1073/pnas.2003968117

Provided by Eindhoven University of Technology

Citation: Velcro method for more precise binding of drug particles (2020, August 24) retrieved 25 July 2024 from <https://phys.org/news/2020-08-velcro-method-precise-drug-particles.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.