

## How biotic and abiotic factors regulate the sediment denitrification rate among wetland types

September 9 2020, by Zhang Nannan



A scaling analysis for estimating nitrate removal in wetlands with varying hydrologic regimes. Credit: MA Lin

Wetlands play a vital role in removing nitrogen (N) from aquatic environments via the denitrification process, which is regulated by multiple environmental and biological factors. Until now, it has been unclear how the denitrification rate responds to abiotic and biotic factors



in wetlands under different hydrological conditions.

Researchers from the Wuhan Botanical Garden of the Chinese Academy of Sciences (CAS) measured the sediment potential denitrification rate (PDR) and unamended denitrification rate (UDR), and quantified denitrifier abundance in 36 stream, river, pond and ditch wetland sites along the Dan River, a nitrogen-rich river in central China.

According to the researchers, sediment denitrification rates significantly differed among streams, rivers, ponds and ditches, with the ditches having the highest denitrification rates.

Meanwhile, denitrification rates showed significant seasonality, and were negatively associated with water velocity and viscosity.

Interestingly, PDR and UDR had positive relationships with N and carbon (C) availability in streams and rivers, but such correlations were not found in ponds and ditches.

Furthermore, the relative nirS gene abundance, total C ratio, Reynolds number and interstitial space jointly provide an estimated nitrate removal in <u>wetlands</u> with varying hydrologic regimes.

These findings highlight that hydrological conditions, especially water velocity and hydrologic pulsing, play a non-negligible role in regulating N biogeochemical processes in wetlands.

This research was supported by the National Natural Science Foundation of China and the Youth Innovation Promotion Association of CAS.

The study is published in *Science of the Total Environment*, titled "Seeking the hotspots of nitrogen removal: A comparison of sediment <u>denitrification</u> rate and denitrifier abundance among wetland types with



different hydrological conditions."

**More information:** Danli Deng et al. Seeking the hotspots of nitrogen removal: A comparison of sediment denitrification rate and denitrifier abundance among wetland types with different hydrological conditions, *Science of The Total Environment* (2020). DOI: 10.1016/j.scitotenv.2020.140253

Provided by Chinese Academy of Sciences

Citation: How biotic and abiotic factors regulate the sediment denitrification rate among wetland types (2020, September 9) retrieved 22 June 2024 from <u>https://phys.org/news/2020-09-biotic-abiotic-factors-sediment-denitrification.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.