

Choanozoan and picozoan marine protists are probably virus eaters: study

September 24 2020




Sampling seawater in the Gulf of Maine. Credit: Ramunas Stepanauskas and coauthors

Viruses occur in astronomic numbers everywhere on Earth, from the atmosphere to the deepest ocean. Surprisingly, considering the abundance and nutrient-richness of viruses, no organisms are known to use them as food. In *Frontiers in Microbiology*, researchers publish the first compelling evidence that two groups of ecologically important marine protists, choanozoans and picozoans, are virus eaters, catching their "prey" through phagocytosis (i.e. engulfing).

"Our data show that many protist cells contain DNA of a wide variety of non-[infectious viruses](#) but not bacteria, strong evidence that they are feeding on viruses rather than on bacteria. That came as a big surprise, as these findings go against the currently predominant views of the role of viruses and protists in the marine food webs," says corresponding author Dr. Ramunas Stepanauskas, Director of the Single Cell Genomics Center at Bigelow Laboratory for Ocean Sciences in East Boothbay, Maine, U.S..

Stepanauskas and colleagues sampled surface seawater from two sites: the Northwestern Atlantic in the Gulf of Maine, U.S. in July 2009, and the Mediterranean off Catalonia, Spain in January and July 2016. They used modern single-cell genomics tools to sequence the total DNA from 1,698 individual protists in the water. Each of the resulting Single Amplified Genomes (SAGs) consists of the genome of a single protist, with or without associated DNA: for example, from symbionts, ingested prey, or viruses or bacteria sticking to its exterior. The technique is very sensitive, but doesn't directly show the type of relationship between a protist and its associates.



Separating protists through flow cytometry. Credit: Ramunas Stepanauskas and coauthors

The researchers found a range of protists including alveolates, stramenopiles, chlorophytes, cercozoans, picozoans, and choanozoans. Nineteen percent of SAGs from the Gulf of Maine and 48% of those from the Mediterranean were associated with bacterial DNA, suggesting that these protists had eaten bacteria. More common were viral sequences, found in 51% of SAGs from the Gulf of Maine and 35% of those from the Mediterranean, with a frequency of 1-52 [virus](#) types per protist. Most were from viruses known to infect bacteria—presumably representing parasites of the protists' bacterial prey.

But choanozoans and picozoans, which only occurred in the Gulf of Maine sample, were different. These groups, neither of which have chloroplasts, are poorly known. Choanozoans (3-10 μm ; also known as choanoflagellates), are of great evolutionary interest as the closest living relatives of animals and fungi. The tiny (up to 3 μ) picozoans were first discovered twenty years ago and originally known as picobiliphytes. Until now, their [food sources](#) were a puzzle, as their feeding apparatus is too small for bacteria—but ample for viruses, most of which are smaller than 150 nm.

Every single one of the choanozoan and picozoan SAGs were associated with [viral sequences](#) from bacteriophages and CRESS-DNA viruses, but mostly without any bacterial DNA, while the same sequences were found across a great diversity of species.



Single-cell genomics laboratory work. Credit: Ramunas Stepanauskas and coauthors

"It is very unlikely that these viruses are capable of infecting all the protists in which they were found," says Dr. Julia Brown, a researcher at the Bigelow Laboratory for Ocean Sciences and coauthor on the study.

The authors conclude that choanozoans and picozoans probably routinely eat viruses.

"Viruses are rich in phosphorus and nitrogen, and could potentially be a good supplement to a carbon-rich diet that might include cellular prey or carbon-rich marine colloids," says Brown. "The removal of viruses from the water may reduce the number of viruses available to infect other organisms, while also shuttling the organic carbon within virus particles higher up the food chain. Future research might consider whether protists that consume [viruses](#) accumulate DNA sequences from their viral prey within their own genomes, or consider how they might protect themselves from infection."

More information: Julia M. Brown et al, Single Cell Genomics Reveals Viruses Consumed by Marine Protists, *Frontiers in Microbiology* (2020). [DOI: 10.3389/fmicb.2020.524828](https://doi.org/10.3389/fmicb.2020.524828)

Provided by Frontiers

Citation: Choanozoan and picozoan marine protists are probably virus eaters: study (2020, September 24) retrieved 22 March 2023 from <https://phys.org/news/2020-09-choanozoan->

picozoan-marine-protists-virus.html

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.