

Deforestation squeezes top predators in forest streams

September 25 2020, by Hayley Dunning



Credit: CC0 Public Domain

Streams in areas converted to oil palm have different top predators to those flowing through intact forests, affecting stream food webs.

Logging and deforestation are known to affect biodiversity and [species richness](#), but a new study shows how these may create trouble for local ecosystems by affecting the [food](#) chain length.

Ecosystems like streams provide valuable [ecosystem services](#), such as provision of fresh water and food and flood prevention, which could be diminished if food chains are altered and the ecosystem fails to function as normal.

A team of researchers led by Imperial College London and the National University of Singapore studied 19 streams in Borneo, in an area with a variety of land uses—primary forest, partially logged areas and full conversion to oil palm plantation—known as the Stability of Altered Forest Ecosystems (SAFE) Project. Their study is published today in the journal *Ecology*.

They found that top predators were disproportionately affected by deforestation, with fewer [predator](#) links in the food chain in oil palm plantation streams. The researchers found that this was caused by a change in the type of predators.

More specialized, endemic predators, such as the barb (*Hampala sabana*) were replaced by generalists, predators such as the non-native snakehead (*Channa striata*) that can eat a wider diet, such as fish, frogs and invertebrates.

Knock-on effects

Lead researcher Dr. Clare Wilkinson, from the University Singapore and formerly of Imperial College London, said: "We know deforestation and land-use changes can affect the [local species](#), but are often unsure what knock-on impacts these changes can have. Our study shows that losses of species richness can affect food chains, and in particular their

length—how many species there are between the primary producers of nutrients at the bottom of the food chain and the top predators. These changes could have knock-on effects for the functioning of the ecosystem."

The team measured carbon and nitrogen in streams and more than 3,000 captured fish, which allowed them to track the sources of nutrients and how these moved through the food chain.

Carbon isotopes—versions of carbon—can reveal where the basic nutrients in an ecosystem come from, either from the stream (by photosynthesising organisms like algae) or from the surrounding environments (such as through leaves and soil entering a stream).

Deforestation could affect both of these sources, as the reduction of forest cover could increase the temperature and sunshine of the water but reduce the amount of material entering a stream.

Preventing further impacts

However, the team found no difference in the amount of carbon from in-stream sources, and no effect of changes in environmental sources of carbon, suggesting these [ecosystems](#) rely more on in-stream sources that remain relatively stable.

The changes in nitrogen, however, revealed that top predators were strongly affected by land-use changes, with different predators found in streams in oil palm plantations.

SAFE Director of Science Professor Rob Ewers, from the Department of Life Sciences at Imperial, said: "The change in ecosystem structure and the loss of specialized species could have knock-on effect on the health of the ecosystem. We now need to do more research to follow the

consequences of these changes, and develop strategies that reverse, mitigate, or prevent further impacts."

More information: Clare L. Wilkinson et al. Forest conversion to oil palm compresses food chain length in tropical streams, *Ecology* (2020). [DOI: 10.1002/ecy.3199](https://doi.org/10.1002/ecy.3199)

Provided by Imperial College London

Citation: Deforestation squeezes top predators in forest streams (2020, September 25) retrieved 2 May 2024 from <https://phys.org/news/2020-09-deforestation-predators-forest-streams.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--