

Researchers discover effective pathway to convert carbon dioxide into ethylene

September 17 2020




Illustration of the ElectroCatalysis system which synthesized the smooth nanowire and then activated it by applying a voltage to get the rough stepped surface that is highly selective for CO_2 reduction to ethylene. Credit: Yu Huang and William A. Goddard III

A research team from Caltech and the UCLA Samueli School of

Engineering has demonstrated a promising way to efficiently convert carbon dioxide into ethylene—an important chemical used to produce plastics, solvents, cosmetics and other important products globally.

The scientists developed nanoscale copper wires with specially shaped surfaces to catalyze a chemical reaction that reduces [greenhouse gas emissions](#) while generating ethylene—a valuable chemical simultaneously. Computational studies of the reaction show the shaped catalyst favors the production of ethylene over hydrogen or methane. A study detailing the advance was published in *Nature Catalysis*.

"We are at the brink of fossil fuel exhaustion, coupled with global climate change challenges," said Yu Huang, the study's co-corresponding author, and professor of materials science and engineering at UCLA.

"Developing materials that can efficiently turn greenhouse gases into value-added fuels and chemical feedstocks is a critical step to mitigate global warming while turning away from extracting increasingly limited fossil fuels. This integrated experiment and [theoretical analysis](#) presents a sustainable path towards [carbon dioxide](#) upcycling and utilization."

Currently, ethylene has a global annual production of 158 million tons. Much of that is turned into polyethylene, which is used in plastic packaging. Ethylene is processed from hydrocarbons, such as natural gas.

"The idea of using copper to catalyze this reaction has been around for a long time, but the key is to accelerate the rate so it is fast enough for industrial production," said William A. Goddard III, the study's co-corresponding author and Caltech's Charles and Mary Ferkel Professor of Chemistry, Materials Science, and Applied Physics. "This study shows a solid path towards that mark, with the potential to transform ethylene production into a greener industry using CO₂ that would otherwise end up in the atmosphere."

Using copper to kick start the carbon dioxide (CO₂) reduction into ethylene reaction (C₂H₄) has suffered two strikes against it. First, the initial chemical reaction also produced hydrogen and methane—both undesirable in industrial production. Second, previous attempts that resulted in ethylene production did not last long, with conversion efficiency tailing off as the system continued to run.

To overcome these two hurdles, the researchers focused on the design of the copper nanowires with highly active "steps"—similar to a set of stairs arranged at atomic scale. One intriguing finding of this collaborative study is that this step pattern across the nanowires' surfaces remained stable under the reaction conditions, contrary to general belief that these high energy features would smooth out. This is the key to both the system's durability and selectivity in producing ethylene, instead of other end products.

The team demonstrated a carbon dioxide-to-[ethylene](#) conversion rate of greater than 70%, much more efficient than previous designs, which yielded at least 10% less under the same conditions. The new system ran for 200 hours, with little change in [conversion efficiency](#), a major advance for copper-based catalysts. In addition, the comprehensive understanding of the structure-function relation illustrated a new perspective to design highly active and durable CO₂ reduction catalyst in action.

Huang and Goddard have been frequent collaborators for many years, with Goddard's research group focusing on the theoretical reasons that underpin [chemical reactions](#), while Huang's group has created new materials and conducted experiments. The lead author on the paper is Chungseok Choi, a graduate student in materials science and engineering at UCLA Samueli and a member of Huang's laboratory.

More information: Chungseok Choi et al, Highly active and stable

stepped Cu surface for enhanced electrochemical CO₂ reduction to C₂H₄, *Nature Catalysis* (2020). [DOI: 10.1038/s41929-020-00504-x](https://doi.org/10.1038/s41929-020-00504-x)

Provided by University of California, Los Angeles

Citation: Researchers discover effective pathway to convert carbon dioxide into ethylene (2020, September 17) retrieved 11 May 2024 from <https://phys.org/news/2020-09-effective-pathway-carbon-dioxide-ethylene.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.