

Electric current is manipulated by light in an organic superconductor



Fig. 1. Spectra of SHG and THG which are emitted by irradiation of 6 fs laser pulses in an organic superconductor, κ -(BEDT-TTF)₂Cu[N(CN)₂]Br. Observation of the SHG in the centrosymmetric materials indicates a non-linear current which cannot be described by Ohm's law. Right panel shows the crystal structure of the sample. Credit: NINS/IMS

A polarized petahertz current is driven by an ultrashort laser in an organic superconductor. This is in contrast to the common sense belief which is justified by Ohm's law, i.e., a net current cannot be induced by an oscillating electric field of light. The current enhances near the superconducting transition temperature. The light-driven petahertz current opens a way to high-speed operation of computers which is one million times faster than conventional ones.



In modern information technology (IT), data are processed and carried by the motions of electrons in a CPU. In the electric circuits, the electrons move in a desired direction by an applied electric field. A frequency of the on-off switching of the electron motion, which is referred to as a "CPU clock" for example, is an order of gigahertz (10^9 Hz).

On the other hand, an oscillating field of <u>light</u> with a frequency of petahertz (10^{15} Hz) has the potential for realizing petahertz operation of the on-off switching. If one can move electrons with the light frequency, the speed of data processing could be one million times faster than those in conventional computers. An electromagnetic oscillation of light has, however, never driven polarized current (i.e., the <u>time</u> average of the current during the light pulse is zero), because the oscillating light field is temporally/spatially symmetric. Researchers at Tohoku University, Nagoya University, Institute for Molecular Science, Okayama Science University and Chuo University have succeeded in moving electrons in an organic superconductor in a specific direction by irradiation of ultrashort laser pulses.



Fig. 2. (a) CEP (carrier-envelope phase) dependence of SHG (b) Temperature dependence of SHG. (Superconducting transition temperature of the sample is



11.5 K.) Credit: NINS/IMS

According to Ohm's law, an induced current (and velocity of electrons) is proportional to the applied electric field. Note that Ohm's law holds, if the electrons are scattered many times in solids. In fact, the resistivity of the materials is determined by the electron-electron and/or electron-phonon scattering processes. If the electric field can be applied on the time scale shorter than the scattering time, however, the electrons in solids do not have enough time to be averaged. Instead, the electrons should be accelerated and generate a polarized net current. Therefore, the researchers have attempted to realize such a 'scattering-free current' using ultrashort laser pulses which are sufficiently shorter than the electron scattering time (about 40 femtoseconds in organic superconductors).

One obstacle to realize such an experiment is that electric detection of such a short-time current is impossible. Therefore, the researchers employ the optical detection. Second harmonic generation (SHG) has been well known as the method for detecting electronic symmetry breaking such as a macroscopic dipole moment in ferroelectrics. The SHG can be also induced by the polarized current which is another type of the electronic symmetry breaking.

The researchers shine their ultrashort laser with a pulse width of about 6 fs $(6 \times 10^{-15} \text{ s})$ on an organic centrosymmetric superconductor, κ -(BEDT-TTF)₂Cu[N(CN)₂]Br, and detect a second harmonic generation (SHG). This is in contrast to the common sense because the SHG is generated only in the materials in which spatial symmetry is broken. Their detection of the SHG in the centrosymmetric material indicates that a polarized net current is generated during the light irradiation.





Fig. 3. Schematic illustration of the SHG induced by a petahertz non-linear current in an organic superconductor κ - BEDT-TTF compounds Credit: NINS/IMS

To confirm such a non-linear polarized current, the researchers investigate carrier-envelope phase (CEP; relative phase between the oscillation of light and its envelope) dependence of the SHG, because CEP sensitive nature is a characteristic behavior of the current induced SHG. A periodic change of the SHG intensity as a function of the CEP is an evidence that the observed SHG is actually attributable to the



scattering-free current.

The researchers further demonstrate that the relation between the scattering-free current and the superconductivity. The present result (Fig. 2b) shows that the SHG is detected at the temperature range below 50 K (> superconducting transition temperature (T_{SC} =11.5 K)). The result also shows the intensity of the SHG rapidly grows toward the transition temperature below 25 K (~2 × T_{SC}), indicating that the scattering-free current is sensitive to a "superconducting fluctuation". In many superconductors, the superconducting fluctuation, or microscopic seeds of superconductivity, has been found at temperatures higher than the superconducting transition, and the increase in the intensity of the second harmonic seems to be related to the superconducting fluctuations.

The researchers say, "With further understanding of the scattering-free non-linear petahertz current, we may be able to make computers with an operation speed of petahertz which is million times faster than the present ones of gigahertz. This phenomenon can be used also as a tool to elucidate the microscopic mechanism of superconducting states, because it is sensitive to the superconducting fluctuation."

More information: Y. Kawakami et al, Petahertz non-linear current in a centrosymmetric organic superconductor, *Nature Communications* (2020). DOI: 10.1038/s41467-020-17776-3

Provided by National Institutes of Natural Sciences

Citation: Electric current is manipulated by light in an organic superconductor (2020, September 4) retrieved 23 June 2024 from <u>https://phys.org/news/2020-09-electric-current-superconductor.html</u>



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.