


Old grasslands show high biodiversity and conservation value

September 10 2020



The number of (a) all, (b) grassland-dependent native and (c) grassland-dependent endangered plant species in each vegetation type. “Old” and “new” refer to old and new grasslands (see detail in the text). Bars and boxes are medians and quartiles, with ranges between minimum and maximum values. An outlier (values less than the first quartile minus 1.5 times the interquartile range [IQR] or greater than the third quartile plus 1.5 times the IQR) is shown by a dot. Credit: University of Tsukuba

"The grass is always greener on the other side," as the saying goes, but in this case, it's more diverse. Researchers from Japan have discovered that

old grasslands have higher plant diversity than new ones, and that grassland longevity can be an indicator of high conservation priority.

In a study published this month in *Ecological Research* online, researchers from the University of Tsukuba have revealed that the longer grasslands have been around, the higher their plant diversity, and the more likely they are to be of high conservation priority.

Grasslands can be classified as natural (existing in natural climatic conditions and disturbance systems) or seminatural (maintained by artificial disturbances such as pasturing, fire or mowing). Seminatural grasslands are ecosystems with rich biodiversity. Unfortunately, both types of grasslands are declining globally.

"There's an urgent need to identify grasslands of high conservation priority," says lead author of the study Taiki Inoue. "The results of a growing number of recent studies show that vegetation history affects current biological communities. The aim of our study was to evaluate whether the uninterrupted continuity of grasslands through time promotes biodiversity, and therefore can be an indicator of conservation priority."

grasslands as refuges for those species."

Old and new grasslands also differed in species composition, with the composition of new grasslands ranging between that of old grasslands and forests. This suggests that new grasslands continue to be affected by past forestation more than 52 years after deforestation. Old grasslands were found to have eleven indicator species, with none found in new grasslands, revealing that the plant community in old grasslands was unique.

"Our findings indicate that grasslands that have been around for a long time are where conservation effort should be focused," says Inoue.

Future studies investigating the effect of vegetation history on the current biodiversity of [grassland](#) plant [species](#) will improve understanding of how biological communities are formed, and will be key to allocating [conservation priority](#).

More information: Taiki Inoue et al. The effects of temporal continuities of grasslands on the diversity and species composition of plants, *Ecological Research* (2020). [DOI: 10.1111/1440-1703.12169](https://doi.org/10.1111/1440-1703.12169)

Provided by University of Tsukuba

Citation: Old grasslands show high biodiversity and conservation value (2020, September 10) retrieved 3 May 2024 from <https://phys.org/news/2020-09-grasslands-high-biodiversity.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.