

Inertial confinement fusion implosions have significant 3-D asymmetries

September 7 2020, by Patricia Koning



Figure a: Model of the capsule in a laser-irradiated hohlraum from view angle (65°, 120°). Typical size and position of diagnostic windows are shown in blue. Figure (b): Calculated reduction of radiation flux on the capsule in a 3-window hohlraum, assuming complete radiation loss through the windows.

Data correlating two factors that lead to implosion asymmetries have brought Lawrence Livermore National Laboratory (LLNL) scientists a step closer to understanding the gap between simulations and



performance of inertial confinement fusion (ICF) experiments at the National Ignition Facility (NIF).

These experiments aim to ignite a propagating fusion burn wave in deuterium-tritium fuel. To achieve ignition, the fuel must reach a selfheating state in which the energy produced exceeds energy loss from expansion, thermal conduction and radiative cooling.

The best-performing ICF implosions, with central hot-spot pressures of 360 Gbar (billion Earth atmospheres) and fusion yields of 50 kilojoules, have begun to show the effects of fusion self-heating. But these experiments still underperform 1 and 2-D simulations, which predict pressures above 500 Gbar and dynamics dominated by self-heating.

By analyzing data from several years of high-yield NIF ICF experiments, researchers have found a correlation between the velocity of the hot-spot implosion and the asymmetry of fuel areal-density (the combined thickness and density of the imploding frozen fusion fuel shell).

The results were reported in a *Physical Review Letters* paper by Laboratory for Laser Energetics physicist Hans Rinderknecht (who performed this work while a Lawrence Fellow at LLNL), LLNL physicist Dan Casey and LLNL colleagues.

"We know that asymmetry is a principal degradation of implosion performance," Casey said. "We found that 3-D asymmetry does exist and is correlated between two key measurements. These results lay the foundation for subsequent work to track down the sources of the asymmetries."

The correlation was consistent across a wide range of shot and target configurations, including shots from the high-density carbon (HDC), "BigFoot" (high-adiabat, or reduced compression, HDC) and traditional



CH (plastic capsule) campaigns.

"We found that most of the implosions performed in the last three years of the ICF program on NIF had an unintentional asymmetry that 'pushed' the implosion to one side instead of imploding it uniformly. This meant that some of the energy was wasted," Rinderknecht said.

The striking coincidence of magnitude and direction between the hotspot velocity and the asymmetry of the fuel areal-density across a wide range of experiments with varied ablator composition, laser-power history and other factors suggests a common, systematic underlying cause. The implosions tended to be pushed toward the diagnostic windows in the hohlraum. Those windows are covered by thinner gold layers that allow X-ray cameras to see the capsule inside the hohlraum.





Neutron-averaged hot-spot flow velocity magnitude and direction evaluated for NIF cryogenic implosions in the HDC (circle), Bigfoot (triangle) and CH (square) campaigns from 2016–2018. The gray symbols represent implosions with bulk velocity below 30 km/s. Credit: Lawrence Livermore National Laboratory

Two overlapping diagnostics

The researchers applied their analysis method to 44 implosions of cryogenic deuterium-tritium ice layers performed on NIF between 2016



and 2018. The hot-spot movement direction was observed in 17 out of 18 HDC, 10 of 11 Bigfoot and six out of 15 CH implosions.

"It's a classic story of how two datasets are better than one," Rinderknecht said.

NADs record how many fusion-produced neutrons make it out of the implosion without scattering. This diagnostic, therefore, can determine the density variations of fuel in various directions.

However, NADs also can be affected by the Doppler shift: if the implosion is moving toward the detector, the neutrons get an energy boost and the signal is therefore increased because of the energy dependent NAD activation cross-section. The nTOF spectrometers also measure the neutron Doppler shift.

Diagnostics are the only way to observe what is actually happening in the implosions, but they can be flawed or misunderstood. "When you start collecting and interpreting data, it's easy to get into arguments about whether what you're seeing is real or not, especially if the result is surprising," Rinderknecht said.

"Once I had both sets of data, I started plotting them together—hot-spot velocity versus density asymmetry—and the shared pattern popped right out," he said. "The two supporting datasets from two different and independently vetted diagnostics confirmed one another across so many shots with such different conditions, it became clear that something real and significant was going on."

Moving forward, LLNL researchers are developing detailed models to assess window radiation losses more quantitatively, including the effects of window architecture and ablation dynamics.



"On the basis of these data a research program has been initiated to find and control the origins of the drive asymmetry, which remains a crucial step for ongoing efforts to achieve ignition on the NIF," the researchers said in the paper.

Preliminary results indicate that asymmetries in laser delivery are comparable to those of the diagnostic windows. Capsule thickness variations, ice-layer thickness variations and laser-to-target misalignment could also be sources of asymmetry.

"Work to identify and control the source of this <u>asymmetry</u> is ongoing and will be essential to further improving implosion performance and achieving ignition in indirect-drive ICF," the researchers said.

Casey and Rinderknecht were joined on the paper, "Azimuthal Drive Asymmetry in Inertial Confinement Fusion Implosions on the National Ignition Facility," by LLNL colleagues Robert Hatarik, Richard Bionta, Brian MacGowan, Prav Patel, Nino Landen, Ed Hartouni and Omar Hurricane.

More information: Hans G. Rinderknecht et al. Azimuthal Drive Asymmetry in Inertial Confinement Fusion Implosions on the National Ignition Facility, *Physical Review Letters* (2020). <u>DOI:</u> <u>10.1103/PhysRevLett.124.145002</u>

Provided by Lawrence Livermore National Laboratory

Citation: Inertial confinement fusion implosions have significant 3-D asymmetries (2020, September 7) retrieved 1 May 2024 from <u>https://phys.org/news/2020-09-inertial-confinement-fusion-implosions-significant.html</u>



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.