

# Research finds mouse populations can be controlled with lower amounts of poison

September 25 2020

---



Credit: Pixabay

Good news for Western Australian wildlife with a recent study suggesting that mice numbers could be controlled with much smaller doses of poison.

ECU Honors student Bridget Duncan tested the resistance of mice

populations in metropolitan Perth to a common and particularly potent brand of rat poison.

Her study showed [house mice](#) are particularly vulnerable to rat poison in Western Australia.

She and her supervisors were surprised to discover local mice lacked a [gene mutation](#) found in mice around the world that helps to resist the effects of rat poison.

"Mice in Europe, the United States and Canada have all been found to carry this mutation and we assumed that mice in Western Australia would be the same," Ms Duncan said.

"However, results told a different story. This is particularly surprising as rat poisons are much more accessible to consumers in Australia compared to other parts of the world and we assumed they would have built up resistance over time."

This was the first investigation to test for this mutation in Australian mice and results may have implications for mouse control around the country.

## **Building resistance over time**

Rat poisons such as warfarin has been used to control rodents for more than 80 years, and as [rats](#) and mice developed resistance to these chemicals, a new generation of more potent chemicals were developed.

"Over time the population gains resistance through animals inheriting the protective genetic mutation from the survivors over generations," Ms Duncan said.

"As a result, the extremely potent second-generation chemicals are now widely used around the world, including Western Australia, where they have a devastating effect on the local wildlife. Small native mammals, birds and reptiles are all killed by these potent poisons, but our research has indicated there might be no need to be so heavy handed with these chemical controls in Western Australia."

Ms Duncan said the research team had speculated why Western Australia mice were so different from populations in other parts of the world, but more research needed to be done.

"Mice in the Perth metropolitan area are all very similar to each other and are most closely related to British population of mice," she said.

"It is possible that Western Australian mice arrived on the early settler ships before this genetic mutation arose in the British population."

Researchers also sampled mice on Browse Island in the remote Kimberley to compare against the Perth metropolitan mice.

"We found the mouse population on Browse Island was more closely related to south-east Asian [mice](#) so probably originated from Indonesian fishing boats," Ms Duncan said.

"They also lack resistance to common poisons so environmental managers can employ a very measured approach to [population](#) control through this method."

Next steps in the research will examine the resistance of rats in Western Australia to the same poisons.

The study "Mus musculus populations in Western Australia lack VKORC1 mutations conferring resistance to first generation

anticoagulant rodenticides: Implications for conservation and biosecurity" can be read in full in the *PLOS ONE* journal.

**More information:** Bridget J.M.L. Duncan et al. *Mus musculus* populations in Western Australia lack VKORC1 mutations conferring resistance to first generation anticoagulant rodenticides: Implications for conservation and biosecurity, *PLOS ONE* (2020). [DOI: 10.1101/2020.07.06.189282](https://doi.org/10.1101/2020.07.06.189282)

Provided by Edith Cowan University

Citation: Research finds mouse populations can be controlled with lower amounts of poison (2020, September 25) retrieved 18 April 2024 from <https://phys.org/news/2020-09-mouse-populations-amounts-poison.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.