

One of the world's driest deserts is the focus of a new study on our changing climate

September 24 2020



University students Ruusa Gottlieb (left) and Priscilla Mundilo measure carbon dioxide release from the soil at one of the high rainfall sites in the Namib Desert. Photo: Heather Throop/ASU

Carbon, one of the main building blocks for all life on Earth, cycles among living organisms and the environment. This cycle, and how it works in one of the driest places on Earth, is the subject of a new study

recently published in the journal *Plant and Soil* with lead author and Arizona State University (ASU) scientist, Heather Throop.

While the [natural carbon cycle](#) should be balanced each year, with about as much [carbon](#) taken out of the atmosphere as is released back by natural processes, humans are upsetting this balance through carbon dioxide additions to the atmosphere, both through changing land use that releases carbon stored in soils and from burning fossil fuels.

In an effort to understand what controls the release of carbon dioxide from soils in deserts, Throop and a team of university students from Namibia conducted [field work](#) in the Namib Desert, one of the world's driest regions that stretches more than 1,200 miles along the Atlantic coasts of Angola, Namibia, and South Africa.

What Throop and her team ultimately determined from their research is that subtle differences in surface topography and erosion have big influences on microorganisms in the soil and these differences ultimately affect carbon cycling. Even in the driest places, they found signs of life influencing carbon cycling.

"The amount of carbon dioxide in the atmosphere affects our climate, so understanding what affects the release of carbon from soils is important predicting how climate will change in the future," says Throop, who is an associate professor in the School of Earth and Space Exploration and the School of Life Sciences.



University students Theopo Iita, Vimbai Marufu, Priscilla Mundilo characterize plant and landscape features at one of the driest sites in the Namib Desert research study. Credit: Throop/ASU

To conduct their analyses, the research team chose six locations that differed in yearly rainfall. At each site they carried out 48-hour sampling campaigns, working continuously day and night to collect data. At each location, the team analyzed the landscape structure and plants and selected representative locations to sample. Then, they simulated rainfall and used gas analyzers to measure carbon dioxide release from soils, to determine how carbon cycling responded as soils dried after the simulated rain.

"It's really an incredible amount of data to collect manually," says Throop. "And having a crew of dedicated and enthusiastic students made this work possible. Often for remote field work like this we just get a snapshot of what is happening at one or two sites or at a few points in time. It was exciting to be able to collect the data continuously for a few days and at six different sites."

The students participating in this research came from the University of Namibia and the Namibia University of Science and Technology. They were each participating in the Summer Drylands Program, an intense research experience where students plan, execute, and report on an experiment within a short timeframe.



University students Rosalia Tshikesho (L) and Etuna Iyambo (R) simulate a rain shower prior to making measurements of soil carbon dioxide release in the

Namib Desert. Credit: Throop/ASU

"The ability of technology to record [soil](#) carbon was outstanding," says co-author and student researcher Vimbai Marufu, who is now in graduate school at the Namibia University of Science and Technology. "What I treasure the most from the experience is what it means to work on an interdisciplinary team and the unexplainable satisfaction of being close to nature."

And there are plans to continue additional fieldwork in the Namib Desert with a recent grant from the National Science Foundation to ASU. This grant will provide support for U.S. students to conduct research in the Namib Desert in collaboration with Namibian researchers. "We hope to use this work to help us in understanding how deserts respond to a changing climate," says Throop. "How biological processes function in the extreme dry of the Namib Desert will give us clues about how relatively wet deserts will behave under drier conditions."

More information: H. L. Throop et al, Multiple scales of spatial heterogeneity control soil respiration responses to precipitation across a dryland rainfall gradient, *Plant and Soil* (2020). [DOI: 10.1007/s11104-020-04614-0](#)

Provided by Arizona State University

Citation: One of the world's driest deserts is the focus of a new study on our changing climate (2020, September 24) retrieved 6 May 2024 from <https://phys.org/news/2020-09-world-driest-focus-climate.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.