

# Forest darkness helps stave off effects of nitrogen pollution – but this is set to change

October 1 2020, by Aisling Irwin

---



The life that grows underfoot accounts for 80% of the biodiversity in a temperate forest. Credit: pikist.com/licenced under CC0

Europe's forests are sitting on a pollution timebomb which could rewrite their ecology when it explodes, say researchers.

Delicate [forest floor](#) plants such as wood sorrel or violet, and the balance among the [tree species](#) that tower above them, are all threatened by decades of accumulated nitrogen pollution. A study has found that the darkness of the forest has subdued the effects of nitrogen. But forests are destined to let in more light in the future as trees succumb to drought and disease.

Forests cover [40% of the European Union's land area](#) and are expanding in some countries, mostly because of active restoration or the abandonment of agricultural land. Forests provide services such as controlling erosion and cycling water but they are also [increasingly threatened](#) by droughts and diseases such as ash dieback.

To understand how they are responding to these challenges it is vital to study the forest floor, says Professor Kris Verheyen, an ecologist at Ghent University in Belgium.

"This herb layer is very often forgotten. Some people call it the step-over layer—you step over it to look at the trees," he said.

Yet, in [temperate forests](#), the life underfoot includes 80% of a forest's biodiversity. The herb layer cycles key nutrients such as phosphorous, potassium and nitrogen, helps decompose tree litter and filters the next generation of trees—since seedlings need to pass through it to embark on their journey to the canopy, says Prof. Verheyen.

## **Records**

In trying to understand more about the forest floor, he led a team that retrieved records, going back in some cases as far as 50 years, of 4,000 forest plots across Europe. For those whose locations were easily identified, the researchers visited to make updated measurements.



The team also carted forest soils from around the continent to their research station where they incorporated them into outdoor experimental environments, called mesocosms, in which they varied the plants' access to nitrogen, temperature and light.

"The basic question we wanted to answer was how do multiple global change drivers determine the trajectories of change over time," says Prof. Verheyen, who led the project, known as [PASTFORWARD](#).



Forest dieback can open up protective canopies. When more light is let in, forests can succumb to drought and disease. Credit: High Contrast/Wikimedia, licenced under CC BY 3.0 DE

Overall, the team found that the key [controlling factor in forest life was light](#), which acts as a bottleneck preventing other changes from exerting an effect.

## **Nitrogen pollution**

One powerful example of this was the way it has held back the effects of nitrogen pollution.

Nitrogen deposition is a chronic problem caused by ammonia emissions from agricultural fertiliser and the creation of nitrogen oxides as a by-product of burning fossil fuels. It draws certain nutrients out of soils, acidifies land, and causes algae to grow in waterways.

The researchers found plenty of nitrogen deposition in forests and documented its consequences for [species](#). But 'the effects are not as strong as we expected because ... the nitrogen is available but the plants (on the forest floor) can't really benefit because they are limited by the amount of light that is available," said Prof. Verheyen.

Some plants—types that tend to be widespread and can survive in a variety of environments including non-[native species](#)—have the machinery to take advantage of an excess of nitrogen and grow more; others—which tend to be specialists with small ranges—do not. In shaded forests they are on an equal footing. But as soon as [the canopy opens up and light pours in](#), those that can exploit the [nitrogen pollution](#) have an advantage.

As a result, European forests are already losing their more specialist species and thus experiencing a drop in biodiversity. Prof. Verheyen is concerned that forest canopies are in danger of opening up as trees die from drought and disease which may open the floodgates to [nitrogen](#).

"That will lead to rapid and very large changes in the herb layer," he said.

Drought—itself a result of climate change—has had a 'massive' effect killing trees in spruce forests in Germany, Belgium and France in recent years although broadleaf forests have been more resistant for now.

This does not mean that broadleaf forests are free from canopy-opening—one example is ash dieback disease. "We do have evidence that because of the dying off of the ash you get a lot of light and then the understorey really explodes because light is no longer a limiting resource," said Prof. Verheyen.

## THE THREE LAYERS OF EUROPEAN TEMPERATE FORESTS



Some experts consider there to be seven layers in a forest, while Prof. Verheyen's team have based their work on three interconnected layers. Credit: Horizon



"These large and probably abrupt changes that may happen in the herb layer will affect the tree regeneration and will certainly determine which species will be able to pass the herb layer filter and which not. It will have its consequences for nutrient cycling because this herb layer really impacts [rate of decomposition](#)."

## Buffering

The researchers also discovered that forests have so far done a remarkable job of buffering plants against the broader climate change going on outside them.

Temperature measurements revealed that forests often have [significantly different temperatures](#) from what weather stations—always placed far from trees—record. In summer, for example, they are on average 4°C cooler. This is not only because thick canopies keep out the light, but also because evapotranspiration of water through the leaves and into the atmosphere sucks heat out of the forest, and the vegetation keeps out breezes that would mix warm air into the cool.

Climate models don't take into account this buffering, despite the fact that two thirds of the world's species live in forests and forest processes such as carbon and nutrient cycling depend on temperature, says Professor Pieter de Frenne, a bioscientist also at Ghent University, who is leading the parallel [FORMICA](#) project investigating forest microclimates.

This, in turn, explains why, in intact forest, there has been [less 'reshuffling' of forest species](#) than was predicted as Europe warms, he says—forest-buffering has allowed many species to cling on.

"We would have expected that forest plants would have responded already to a stronger degree—so that more warm-adapted species would have come into the community and more cold affinity species would have declined or even become locally extinct."

But the effect can't last forever and if the canopy is opened up these species will have a rude awakening as their world warms up to the temperature outside the forest.

"Buffering is buying us time so species get a chance to adapt to the new climate," he said.

The work has practical implications for the way forests are managed and the [PASTFORWARD](#) and [FORMICA](#) teams are now hoping to develop a tool to help [forest](#) managers work out how much of the canopy they can remove—for example for harvesting or as part of the cycle of tree-thinning known as coppicing—without triggering this explosive growth.

Provided by Horizon: The EU Research & Innovation Magazine

Citation: Forest darkness helps stave off effects of nitrogen pollution – but this is set to change (2020, October 1) retrieved 11 May 2024 from <https://phys.org/news/2020-10-forest-darkness-stave-effects-nitrogen.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--