

Genome and satellite technology reveal recovery rates and impacts of climate change on southern right whales

October 29 2020, by Emma Carroll



Credit: University of Auckland tohorā research team, Department of Conservation permit DJI

After close to a decade of globe-spanning effort, the genome of the southern right whale has been <u>released this week</u>, giving us deeper insights into the histories and recovery of whale populations across the southern hemisphere.



Up to 150,000 <u>southern right whales</u> were killed between 1790 and 1980. This whaling drove the <u>global population</u> from perhaps 100,000 to as few as 500 <u>whales</u> in 1920. A century on, we estimate there are 12,000 southern right whales globally. It's a remarkable conservation success story, but one facing new challenges.

The genome represents a record of the different impacts a species has faced. With statistical models we can use genomic information to reconstruct historical <u>population</u> trajectories and patterns of how species interacted and diverged.

We can then link that information with historical habitat and climate patterns. This look back into the past provides insights into how species might respond to future changes. Work on <u>penguins</u> and <u>polar bears</u> has already shown this.

But we also have a new and surprising short-term perspective on the population of whales breeding in the subantarctic Auckland Islands group—Maungahuka, 450km south of New Zealand.

Spying on whales via satellite

Known as tohorā in New Zealand, southern right whales once wintered in the bays and inlets of the North and South Islands of Aotearoa, where they gave birth and socialised. Today, the main nursery ground for this population is Port Ross, in the subantarctic Auckland Islands.

Adult whales socialise at both the Auckland and Campbell Islands during the austral winter. Together these subantarctic <u>islands</u> are internationally recognised as an <u>important marine mammal area</u>.





A southern right whale calf breaches in the subantarctic Auckland Islands. Credit: University of Auckland tohorā research team, Author provided

In August 2020, I led a <u>University of Auckland</u> and <u>Cawthron Institute</u> expedition to the Auckland Islands. We collected small skin samples for genetic and <u>chemical analysis</u> and <u>placed satellite tags on six tohorā</u>. These tags allowed us to follow their migrations to offshore feeding grounds.

It matters where tohorā feed and how their populations recover from whaling because the species is recognised as a sentinel for climate change throughout the Southern Hemisphere. They are what we describe as "capital" breeders—they fast during the breeding season in wintering grounds like the Auckland Islands, living off fat reserves gained in



offshore feeding grounds.

Females need a lot in the "bank" because their calves need a lot of energy. At 4-5m at birth, these calves can grow up to <u>a metre a month</u>. This investment costs the <u>mother 25% of her size</u> over the first few months of her calf's life. It's no surprise that calf growth depends on the <u>mother being in good condition</u>.

Females can only breed again once they've regained their fat capital. Studies in the South Atlantic show wintering grounds in <u>Brazil</u> and <u>Argentina</u> produce more calves when prey is more abundant, or environmental conditions suggest it should be.

The first step in understanding the relationship between recovery and prey in New Zealand is to identify where and on what tohorā feed. The potential feeding areas for our New Zealand population could cover roughly a third of the Southern Ocean. That's why we turn to technologies like satellite tags to help us understand where the whales are going and how they get there.

Where tohorā go

So far, all tracked whales have migrated west; away from the historical whaling grounds to the <u>east near the Chatham Islands</u>. As they left the Auckland Islands, two whales visited other oceanic islands—skirting around Macquarie Island and visiting Campbell Island.

It also seems one whale (Bill or Wiremu, identified as male using genetic analysis of his skin sample) may have reached his feeding grounds, likely at the subtropical convergence. The clue is in the pattern of his tracks: rather than the continuous straight line of a whale migrating, it shows the doughnuts of a whale that has found a prey patch.





Migratory track of southern right whale Bill/Wiremu, where the convoluted track could indicate foraging behaviour.

The subtropical convergence is an area of the ocean where temperature



and salinity can change rapidly, and this can aggregate whale prey. Two whales we tracked offshore from the Auckland Islands in 2009 <u>visited</u> <u>the subtropical convergence</u>, but hundreds of kilometres to the east of Bill's current location.

As Bill and his compatriots migrate, we've begun analysing data that will tell us about the recovery of tohora in the past decade. The most recent population size estimate we have is from 2009, when there were about 2,000 whales.

I am using genomic markers to learn about the kin relationships and, in doing so, the population's size and growth rate. Think of it like this. Everybody has two parents and if you have a small population, say a small town, you are more likely to find those parents than if you have a big population, say a city.

This nifty statistical trick is known as the "<u>close kin</u>" approach to estimating population size. It relies on detailed understanding of the kin relationships of the whales—something we have only really been able to do recently using new genomic sequencing technology.

Global effort to understand climate change impacts

Globally, southern right whales in <u>South Africa</u> and Argentina have bred less often over the past decade, leading to a <u>lower population growth rate</u> in Argentina.

Concern over this slowdown in recovery has prompted researchers from around the world to work together to understand the relationship between climate change, foraging ecology and recovery of southern right whales as part of the International Whaling Commission <u>Southern Ocean</u> <u>Research Partnership</u>.



The genome helps by giving us that long view of how the whales responded to climate fluctuations in the past, while satellite tracking gives us the short view of how they are responding on a day-to-day basis. Both will help us understand the future of these amazing creatures.

This article is republished from <u>The Conversation</u> under a Creative Commons license. Read the <u>original article</u>.

Provided by The Conversation

Citation: Genome and satellite technology reveal recovery rates and impacts of climate change on southern right whales (2020, October 29) retrieved 24 May 2024 from <u>https://phys.org/news/2020-10-genome-satellite-technology-reveal-recovery.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.