

## Microscopy breakthrough reveals how proteins behave in 3-D

October 22 2020



A new microscopy system that can can image individual molecules in 3D and capture the way they "wobble" uses a specially engineered glass plate developed by University of Rochester optical scientists. Credit: University of Rochester photo / J. Adam Fenster



Six years ago, the Nobel Prize in chemistry was awarded to three scientists for finding ways to visualize the pathways of individual molecules inside living cells.

Now, researchers at the University of Rochester and the Fresnel Institute in France have found a way to visualize those molecules in even greater detail, showing their position and orientation in 3-D, and even how they wobble and oscillate. The work could shed invaluable insights into the <u>biological processes</u> involved, for example, when a cell and the proteins that regulate its functions react to the virus that causes COVID-19.

"When a protein changes shape, it exposes other atoms that enhance the biological process, so the change of shape of a protein has a huge effect on other processes inside the cell," says Sophie Brasselet, director of the Fresnel Institute, who collaborated with Miguel Alonso and Thomas Brown, both professors of optics at Rochester.

Nicknamed CHIDO—for "Coordinate and Height super-resolution Imaging with Dithering and Orientation"—the technology is described in a new paper published in *Nature Communications*. Designed and built by lead authors Valentina Curcio, a Ph.D. student in Brasselet's group, and Luis Aleman-Castaneda, a Ph.D. student in Alonso's group, CHIDO is precise within "tens of nanometers in position and a few degrees of orientation" in determining the parameters of single molecules," the team reports.

Using a <u>glass plate</u> subjected to uniform stress all around its periphery, the device can create and extrapolate wavelength oscillations and changes in polarization that occur when molecules are observed in a fluorescence microscope. The new technology transforms the image of a single molecule into a distorted focal spot, the shape of which directly encodes more precise 3-D information than previous measurement tools. In effect, CHIDO can produce beams that have every possible



polarization state.

"This is one of the beauties of optics," Brown says. "If you have a device that can create just about any polarization state, then you also have a device that can analyze just about any possible polarization state."

The glass plate originated in Brown's lab as part of his long interest in developing beams with unusual polarizations. Alonso, an expert on the theory of polarization, worked with Brown on ways to refine this "very simple but very elegant device" and expand its applications. During a visit to Marseille, Alonso described the plate to Brasselet, an expert in novel instrumentation for fluorescence and nonlinear imaging. Brasselet immediately suggested its possible use in the microscopy techniques she was working on to image individual molecules.

"It's been a very complementary team," Brasselet says.

## 20 years in the making

In 1873, Ernst Abbe stipulated that microscopes would never obtain better resolution than half the wavelength of light. That barrier stood until Nobel laureates Eric Betzig and William Moerner—with their single-molecule microscopy—and Stefan Hell—with his stimulated emission depletion microscopy—found ways to bypass it.





Thomas Brown, professor of optics, holds a plate of glass that has been engineered for a new microscopy system that can image individual molecules in three dimensions as well as capture how they "wobble." Credit: University of Rochester photo / J. Adam Fenster

"Due to their achievements the optical microscope can now peer into the nanoworld," the Nobel committee reported in 2014.

"What was missing in that Nobel Prize and the work in subsequent years was the ability to not only accurately know the location of a molecule, but to be able to characterize its direction and especially its motion in



three dimensions," Brown says.

In fact, the solution Brown, Alonso, and Brasselet now describe had its origins 20 years ago.

Starting in 1999, Brown and one of his Ph.D. students, Kathleen Youngworth, began investigating unusual optical beams that displayed unusual patterns of optical polarization, the orientation of the optical wave. Some of the beams exhibited a spoke-like radial pattern with intriguing properties.

Youngworth demonstrated on a tabletop that, when tightly focused, the beams exhibited polarization components that pointed in almost any direction in three dimensions.

Alexis Spilman Vogt, another Ph.D. candidate, then worked with Brown on creating the same effects by applying stress to the edges of a glass cylinder. Brown's brother-in-law, Robert Sampson, a skilled tool and die specialist, was called upon to fabricate some samples and fit them in metal rings for use with a confocal microscope.

This involved heating both the glass and metal rings. "Metal expands at a faster rate when you heat it than glass does," Brown says, "and so you could heat the glass and metal up very hot, insert the glass in the middle of the metal, and as it cools down the metal would shrink and create a tremendous force on the periphery of the glass."

Sampson inadvertently applied more stress than called for with one of the plates. As soon as his brother-in-law handed it to him, Brown knew the plate had unusual qualities. The Rochester group introduced the term "stress engineered optic" to describe the elements and, as they learned more about both the physical behavior and the mathematics, they realized that the windows could be the path the solving entirely new



problems in microscopy.

And that was the origin of what is now CHIDO, which, coincidently, happens to be Mexican slang for "cool."

"At the time Alexis and I knew the stress-engineered glass was interesting, and would likely have useful applications; we just didn't know at the time what they might be," Brown says. Now, thanks to his collaboration with Alonso and Brasselet, he hopes CHIDO will "catch the imagination" of other researchers in the field who can help further refine and apply the technology.

**More information:** Valentina Curcio et al. Birefringent Fourier filtering for single molecule coordinate and height super-resolution imaging with dithering and orientation, *Nature Communications* (2020). DOI: 10.1038/s41467-020-19064-6

Provided by University of Rochester

Citation: Microscopy breakthrough reveals how proteins behave in 3-D (2020, October 22) retrieved 21 May 2024 from <u>https://phys.org/news/2020-10-microscopy-breakthrough-reveals-proteins-d.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.