


Lighting the path to recycling carbon dioxide

October 6 2020



SEM image shows the dense and uniform cathodic biofilm, which mainly comprises chemolithoautotrophs, and could serve as biocatalysts for efficient carbon dioxide conversion to acetate. Credit: 2020 KAUST

Semiconductive photocatalysts that efficiently absorb solar energy could reduce the energy required to drive a bioelectrochemical process that converts CO₂ emissions into valuable chemicals, KAUST researchers have shown.

Recycling CO₂ could simultaneously reduce [carbon emissions](#) into the atmosphere while generating useful chemicals and fuels, explains Bin

Bian, a Ph.D. student in Pascal Saikaly's lab, who led the research. "Microbial electrosynthesis (MES), coupled with a renewable [energy](#) supply, could be one such technology," Bian says.

MES exploits the capacity of some microbes to take up CO₂ and convert it into chemicals, such as acetate. In nature, chemolithoautotroph microbes metabolize minerals as a source of energy in a process that involves the shuttling of electrons. This capability can be exploited to turn CO₂ into value-added products if the microbes are supplied with a stream of electrons and protons from anodic water splitting in an electrochemical cell (see image).

In their latest work, rather than focus on the CO₂-to-acetate step, the team worked on reducing the energy input for [molecular oxygen](#) (O₂) production at the anode, a reaction that keeps the overall cell in balance. "In MES systems, the process that consumes the most energy is believed to be the oxygen evolution reaction (OER)," Bian explains. Researchers have used light-capturing anode materials, such as [titanium dioxide](#), that harness energy from sunlight to help drive the OER. In their current work, the team investigated a promising alternative for the photoanode, the light-harvesting material, bismuth vanadate.

Bismuth vanadate absorbed energy from a much broader range of the solar spectrum than titanium dioxide, making the whole MES cell more efficient, the team showed. "We obtained solar-to-acetate conversion efficiency of 1.65 percent, which is the highest reported so far," Saikaly says. "This efficiency is around eight times higher than the 0.2 percent efficiency of global natural photosynthesis, which is nature's solar-powered process for converting CO₂ into energy-rich molecules," Bian notes.

So far the team has kept the microbe biocatalysts supplied with a steady stream of electrons and CO₂ to sustain their growth. "The next step for

us is to test our system under real sunlight and monitor the resilience of the biocatalysts under an intermittent renewable energy source," Saikaly says.

More information: Bin Bian et al, Efficient solar-to-acetate conversion from CO₂ through microbial electrosynthesis coupled with stable photoanode, *Applied Energy* (2020). [DOI: 10.1016/j.apenergy.2020.115684](https://doi.org/10.1016/j.apenergy.2020.115684)

Provided by King Abdullah University of Science and Technology

Citation: Lighting the path to recycling carbon dioxide (2020, October 6) retrieved 1 May 2024 from <https://phys.org/news/2020-10-path-recycling-carbon-dioxide.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--