


Physicists successfully carry out controlled transport of stored light

October 13 2020



For the experiment, atoms of rubidium-87 are first pre-cooled and then transported to the main test area, which is a custom-made vacuum chamber. There they are cooled to temperatures of just a few microkelvins. Credit: Windpassinger group

A team of physicists led by Professor Patrick Windpassinger at Johannes Gutenberg University Mainz (JGU) has successfully transported light stored in a quantum memory over a distance of 1.2 millimeters. They have demonstrated that the controlled transport process and its dynamics has only little impact on the properties of the stored light. The researchers used ultra-cold rubidium-87 atoms as a storage medium for the light as to achieve a high level of storage efficiency and a long lifetime.

"We stored the light by putting it in a suitcase so to speak, only that in our case the suitcase was made of a cloud of cold atoms. We moved this suitcase over a short distance and then took the light out again. This is very interesting not only for physics in general, but also for [quantum communication](#), because light is not very easy to 'capture', and if you want to transport it elsewhere in a controlled manner, it usually ends up being lost," said Professor Patrick Windpassinger, explaining the complicated process.

The controlled manipulation and storage of quantum information as well as the ability to retrieve it are essential prerequisites for achieving advances in quantum communication and for performing corresponding computer operations in the quantum world. Optical quantum memories, which allow for the storage and on-demand retrieval of quantum information carried by light, are essential for scalable quantum communication networks. For instance, they can represent important building blocks of quantum repeaters or tools in linear quantum computing. In recent years, ensembles of atoms have proven to be media well suited for storing and retrieving optical quantum information. Using a technique known as electromagnetically induced transparency (EIT), incident light pulses can be trapped and coherently mapped to create a collective excitation of the [storage](#) atoms. Since the process is largely reversible, the light can then be retrieved again with high efficiency.

The future objective is to develop a racetrack memory for light

In their recent publication, Professor Patrick Windpassinger and his colleagues have described the actively controlled transport of such stored light over distances larger than the size of the [storage medium](#). Some time ago, they developed a technique that allows ensembles of cold atoms to be transported on an 'optical conveyor belt' which is produced by two laser beams. The advantage of this method is that a relatively large number of atoms can be transported and positioned with a high degree of accuracy without significant loss of atoms and without the [atoms](#) being unintentionally heated. The physicists have now succeeded in using this method to transport atomic clouds that serve as a light [memory](#). The stored information can then be retrieved elsewhere. Refining this concept, the development of novel quantum devices, such as a racetrack memory for [light](#) with separate reading and writing sections, could be possible in the future.

More information: Wei Li et al, Controlled Transport of Stored Light, *Physical Review Letters* (2020). [DOI: 10.1103/PhysRevLett.125.150501](https://doi.org/10.1103/PhysRevLett.125.150501)

Provided by Universitaet Mainz

Citation: Physicists successfully carry out controlled transport of stored light (2020, October 13) retrieved 1 May 2024 from <https://phys.org/news/2020-10-physicists-successfully.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.