

Probing water for an electrifying cause

October 29 2020



KAUST researchers observed that "pendant" droplets formed from hydrophobic capillaries to uniform electric fields. Credit: KAUST; Anastasia Serin

An experiment, elegant in its simplicity, helps explain why water becomes electrified when it touches hydrophobic surfaces.

For over a century, scientists have been puzzled by the electrification of <u>water</u> when it is brought in contact with water-repellent or 'hydrophobic' materials, such as paraffin wax, oils, air bubbles and perfluorinated membranes and sheets. Underlying mechanisms remain hotly debated. Now, a team of KAUST engineers has untangled the roles of water,



hydrophobicity and <u>environmental factors</u> in this process. This fundamental contribution could support development of better devices for microfluidics and nanofluidics and for generating clean energy.

"Hydrophobic surfaces are quite common," notes Jamilya Nauruzbayeva, Ph.D. student and lead author of the study. "For instance, polypropylene and perfluorinated pipettes, tubes, coatings and membranes are <u>hydrophobic surfaces</u> used for many basic sciences and engineering applications. Thus, it is important to understand which mechanisms are at play to improve them and develop new ones."

Himanshu Mishra, who conceived and led this study, says that he has been thinking about this problem for over five years. "Probing the <u>surface</u> of water is an excruciatingly difficult undertaking because the thickness of interfaces is down to the molecular scale, which no experimental techniques can probe unambiguously," Mishra explains.

"This is an electrifying subject at water conferences; over the years, through experiment and theory, several competing factors and mechanisms have been proposed," says Mishra. These include, for instance, the dipolar nature of the water molecule; the instantaneous charge transfer between interfacial water molecules and hydrophobes; the dissolution of atmospheric CO2in water; and the interfacial accumulation of intrinsic ions of water (i.e., hydroxide and hydronium ions).

Mishra and his students teamed up with Carlos Santamarina to design elemental experiments to unentangle the role of water, its ions and pH, hydrophobicity of surfaces, and environmental factors, such as relative humidity and CO2content.

Using a parallel plate capacitor, they exposed "pendant" droplets formed from hydrophobic capillaries respond to uniform electric fields. The



competition between their weight and the electrical force tilted the pendant droplets, which revealed their charge.



Hydrophobic surfaces (1) are intrinsically negatively charged and so when the hydrophobic capillary (2) draws in water from the (neutral) water reservoir, it selectively attracts cations (i.e., positively charged ions) that populate the electrical double layer (shown in the insert). Thus, when the capillary (3) is pulled out of the water reservoir, it carries water with a net-positive charge leaving behind an equal and opposite charge in the water reservoir. When the water is dispensed (4), hydrophobicity ensures that the entire volume of water is expelled with the excess positive charge. Credit: Nauruzbayeva et al.; KAUST; Ivan Gromicho

Next, they utilized an electrometer—capable of measuring charges down to a few electrons—to measure the charge of the water reservoirs from where the droplets were withdrawn. They discovered that as a water droplet is withdrawn using a hydrophobic capillary, the water reservoir acquires an equal and opposite negative charge. This is not the case when you use a glass capillary.

"From these experimental results, we could deduce that these



hydrophobic surfaces carried negative surface charge, even in air, which is quite counterintuitive," explains Nauruzbayeva. "When the surface is inserted in water, positive ions are attracted toward it and negative ions are repelled. Hydrophobicity ensures that the liquid departs from the surface without leaving a film behind."

"This discovery was born out of a deep understanding of the science concepts coupled with simple scientific elegance," says Santamarina. Mishra agrees by concluding that "the strength of our contribution lies in its simplicity."

More information: Jamilya Nauruzbayeva et al. Electrification at water–hydrophobe interfaces, *Nature Communications* (2020). DOI: 10.1038/s41467-020-19054-8

Provided by King Abdullah University of Science and Technology

Citation: Probing water for an electrifying cause (2020, October 29) retrieved 24 May 2024 from <u>https://phys.org/news/2020-10-probing-electrifying.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.