

# Predicting tornadoes on UK cold fronts for the first time

October 20 2020

---



Tornado damage on 17 November 2016, near Crewgreen, Shropshire Credit: Matthew Clark

Weather forecasters can more accurately predict when a tornado is likely to hit the UK thanks to a new tool devised in a partnership between the

University of Leeds and the Met Office.

Around 30 [tornadoes](#) occur in the UK each year, 40% of which develop on [cold fronts](#)—but a lack of forecasting methods for these conditions means they strike without warning.

Now researchers at Leeds and the Met Office have for the first time created a prediction for how likely tornadoes are to occur on cold fronts, meaning a more accurate assessment of tornado risk can be made before a cold front crosses the UK.

Matthew Clark, a Met Office scientist who is currently studying for a Ph.D. at Leeds's School of Earth and Environment, said: "Tornadoes are a relatively common [weather](#) hazard on UK cold fronts, but the Met Office have never before had any way of predicting which cold fronts are likely to produce tornadoes, nor did we understand why tornadoes occurred in some fronts but not others.

"These findings should help to improve the UK forecasts of localized, intense wind damage associated with these kinds of weather system. This should enable organizations and people to take precautions and minimize damage and risk."



Damage caused by a tornado which struck on 17 November, 2016, in Ternhill, Shropshire Credit: Matthew Clark

Mr Clark and Douglas Parker, Professor of Meteorology at Leeds' School of Earth and Environment, analyzed tornado reports from the Tornado and Storm Research Organisation (TORRO), radar imagery and surface analysis charts from 114 weather events over a 35 year period. One such event was the largest tornado outbreak in European history when 104 tornadoes touched down across England and Wales on 23 November, 1981, leaving a trail of damage in their wake.

The researchers identified patterns across the weather events, establishing which cold fronts were likely to produce a single tornado, which could produce several, and which would produce none. The research found that most of these tornadoes form when a region of strong winds approaches the front from the cold side. This creates a bulge in the front which helps to make it sharper, increasing the contrast



in wind speed and direction across the front. Where this contrast increases over time, tornadoes are more likely to "spin up" along the front. Occasionally, relatively large outbreaks of tornadoes can occur in this situation. Forecasters can recognize these weather patterns, which alert them to the general risk of a tornado.

Clark and Parker also used their findings to create a predictive tool, which is already being used on an [experimental basis](#) by the Met Office to pinpoint regions at increased risk of tornadoes. Using the wind fields ahead and behind the cold front, forecasters can now compute a percentage probability that tornadoes will happen.

The tool is currently being tested in the Met Office, and was put to use on 29 February this year, successfully predicting the risk of tornadoes in southeast England, with a tornado occurring in Kent as the cold front swept through during the morning. Although use of the product is experimental at the moment, it is hoped that in time it will allow for appropriate messaging to be issued to local responders and other organizations such as airports.

Met Office Chief Meteorologist Paul Davies said: "By explaining, for the first time, "how" and "why" tornadoes form in a given weather system, meteorologists are much better prepared to anticipate events in advance, to interpret and challenge the results of numerical weather prediction models, and to communicate their confidence in a given forecast."

**More information:** Matthew R. Clark et al, Synoptic-scale and mesoscale controls for tornadogenesis on cold fronts: A generalised measure of tornado risk and identification of synoptic types, *Quarterly Journal of the Royal Meteorological Society* (2020). [DOI: 10.1002/qj.3898](https://doi.org/10.1002/qj.3898)

Provided by University of Leeds

Citation: Predicting tornadoes on UK cold fronts for the first time (2020, October 20) retrieved 13 March 2024 from <https://phys.org/news/2020-10-tornadoes-uk-cold-fronts.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.