

Emerald ash borer puts trees on path to functional extinction

November 6 2020, by Brian Wallheimer



Ash trees, like the one seen here, are weakened as emerald ash borer larvae feed under their bark. The invasive pests eventually kill the trees and have been spreading throughout the United States for nearly two decades. Purdue's Songlin Fei has shown that ash trees are not regenerating well in forests where the ash borers have invaded, leaving the tree species likely to become functionally extinct. Credit: Songlin Fei

Since the emerald ash borer's introduction to the United States at the beginning of the 21st century, forest ecologists and government officials have striven to stem its destruction of ash forests. Despite those efforts, the invasive pest may be winning the war.

Mining 16 years of U.S. Forestry Service Forest Inventory Analysis data for 960 counties, Purdue University professor Songlin Fei has shown that in impacted areas, [young trees](#) are dying before they can reach their reproductive stages. Unable to compete with larger trees or resist the [emerald ash borer](#), American ash trees may be doomed to functional extinction.

"As we see overstory ash trees dying in these forests, we had hoped that ash would respond to the emerald ash borer like many forests do to fire, regenerating after the fire is gone," said Fei, a professor of forestry and natural resources and Purdue Agriculture's Dean's Chair of Remote Sensing. "What we found is that these ash trees are not surviving. There is more mortality in the small-diameter ash trees than there is recruitment, and few are maturing into large-diameter trees capable of producing seed. In the long run, we will run out of material and no longer have ash reaching the reproductive age."

Emerald ash borer, a beetle native to northeast Asia, was first detected in Michigan in 2002. Its western range has reached South Dakota in the

north down to Texas in the south. It's in every state east of that line except for Mississippi and Florida. Its larvae burrow under ash tree bark to feed, eventually weakening and killing trees, with losses estimated in the tens of millions so far.

Fei and colleagues from the U.S. Department of Agriculture Forest Service examined records from the U.S. Department of Agriculture's Animal and Plant Health Inspection Service (APHIS) to catalog the locations of emerald ash borer infestations from three periods: 2002–06, 2007–12, and 2013–18. They paired that with the Forest Inventory Analysis data on tree populations collected from 2002-2007 and 2013-2018 to see how the invasions affected ash trees at different points after the ash borer first appeared in an area.

They discovered that in the years near emerald ash borer's introduction, ash seedling populations were abundant in the forests but started to thin out without growing to maturity. In later years, the density of young trees declined significantly. Later periods had 500 fewer seedlings per hectare and sapling recruitment was 50% lower than the tree mortality rate.

"Despite this widespread regeneration, ash trees in the smallest overstory class died at faster rates than they were recruited from seedlings or saplings, resulting in negative population trajectories on plots that have been invaded for more than 10 years," according to the authors' findings, which will be published in the January issue of *Forest Ecology and Management*. "This trend suggests that ash will continue to decline in abundance and may become functionally extinct across the invaded range of emerald ash borer."

Fei cautioned that the results do not suggest that ash trees will disappear altogether from the U.S. However, being functionally extinct means that they will not provide economic value as hardwoods or the ecological services they long have in forested areas.

"Like elm, which was decimated by the Dutch elm disease, there may be places with ash trees and seedlings in the understory, but they won't mature in populations large enough to fulfill their traditional economic or ecological purposes," Fei said. "There also are peripheral areas that haven't been invaded by the emerald ash borer yet. In those places, we still have a chance and we do want to fight there to keep those forests safe."

Ash [trees](#) in residential areas can be treated with insecticides to keep emerald ash borers away. People are also encouraged to buy firewood locally and not to move it long distances that could help the pest expand its range.

More information: Emerald Ash Borer Management:
extension.entm.purdue.edu/EAB/Management.html

Provided by Purdue University

Citation: Emerald ash borer puts trees on path to functional extinction (2020, November 6)
retrieved 4 May 2024 from <https://phys.org/news/2020-11-emerald-ash-borer-trees-path.html>

| |
|--|
| <p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p> |
|--|