

Offshore submarine freshwater discovery raises hopes for islands worldwide

November 25 2020




Eric Attias (middle) deploying the surface-towed marine controlled-source electromagnetic (CSEM) system off of Hawai'i Island. Credit: University of Hawai'i

Twice as much freshwater is stored offshore of Hawai'i Island than was previously thought, according to a University of Hawai'i study with important implications for volcanic islands around the world. An extensive reservoir of freshwater within the submarine southern flank of the Hualālai aquifer has been mapped by UH researchers with the Hawai'i EPSCoR 'Ike Wai project. The groundbreaking findings, published in *Science Advances*, reveal a novel way in which substantial volumes of freshwater are transported from onshore to offshore submarine aquifers along the coast of Hawai'i Island.

This mechanism may provide alternative renewable resources of freshwater to volcanic islands worldwide. "Their evidence for separate freshwater lenses, stacked one above the other, near the Kona coast of Hawai'i, profoundly improves the prospects for [sustainable development](#) on volcanic islands," said UH Mānoa School of Ocean and Earth Science and Technology (SOEST) Dean Brian Taylor.

Paradigm shift

Through the use of marine controlled-source electromagnetic imaging, the study revealed the onshore-to-offshore movement of freshwater through a multilayer formation of basalts embedded between layers of ash and soil, diverging from previous groundwater models of this area. Conducted as a part of the National Science Foundation-supported 'Ike Wai project, research affiliate faculty Eric Attias led the marine geophysics campaign.




Porpoise system graphic. Credit: University of Hawai'i

"Our findings provide a paradigm shift from the conventional hydrologic conceptual models that have been vastly used by multiple studies and water organizations in Hawai'i and other volcanic islands to calculate sustainable yields and aquifer storage for the past 30 years," said Attias. "We hope that our discovery will enhance future hydrologic models, and consequently, the availability of clean freshwater in volcanic islands."

Co-author Steven Constable, a professor of geophysics at the Scripps Institution of Oceanography, who developed the controlled source electromagnetic system used in the project, said, "I have spent my entire career developing marine electromagnetic methods such as the one used here. It is really gratifying to see the equipment being used for such an impactful and important application. Electrical methods have long been used to study groundwater on land, and so it makes sense to extend the application offshore."

Kerry Key, an associate professor at Columbia University who employs electromagnetic methods to image various oceanic Earth structures, who not involved in this study, said, "This new electromagnetic technique is a game changing tool for cost-effective reconnaissance surveys to identify

regions containing [freshwater aquifers](#), prior to more expensive drilling efforts to directly sample the pore waters. It can also be used to map the lateral extent of any aquifers already identified in isolated boreholes."



Credit: University of Hawai'i

Two-times more water

Donald Thomas, a geochemist with the Hawai'i Institute of Geophysics and Planetology in SOEST who also worked on the study, said the findings confirm two-times the presence of much larger quantities of stored groundwater than previously thought.

"Understanding this new mechanism for groundwater...is important to better manage groundwater resources in Hawai'i," said Thomas, who leads the Humuula Groundwater Research project, which found another large freshwater supply on Hawai'i Island several years ago.



University of Hawai'i Research Affiliate Faculty Eric Attias at Wailupe Beach Park on O'ahu. Credit: University of Hawai'i

Offshore freshwater systems similar to those flanking the Hualālai aquifer are suggested to be present for the island of O'ahu, where the electromagnetic imaging technique has not yet been applied, but, if demonstrated, could provide an overall new concept to manage freshwater resources.

The study proposes that this newly discovered transport mechanism may

be the governing mechanism in other volcanic [islands](#). With offshore reservoirs considered more resilient to [climate change](#)-driven droughts, [volcanic islands](#) worldwide can potentially consider these resources in their water management strategies.

More information: E. Attias et al., "Marine electrical imaging reveals novel freshwater transport mechanism in Hawai'i," *Science Advances* (2020). [advances.sciencemag.org/lookup ... 1126/sciadv.abd4866](https://advances.sciencemag.org/lookup...1126/sciadv.abd4866)

Provided by University of Hawaii at Manoa

Citation: Offshore submarine freshwater discovery raises hopes for islands worldwide (2020, November 25) retrieved 20 March 2024 from <https://phys.org/news/2020-11-offshore-submarine-freshwater-discovery-islands.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
