

Investigating optical activity under an external magnetic field

November 6 2020



Credit: Petr Kratochvil/public domain

Optical activity in chiral molecules has become a hot topic in physics and optics, representing the ability to manipulate the polarized state of light. Understanding how molecules rotate the plane of plane-polarized light has widespread applications, from analytic chemistry to biology and medicine—where it can, for example, be used to detect the amount of sugar in a substance. A new study published in *EPJ B* by Chengping Yin

of the Guangdong Provincial Key Laboratory of Quantum Engineering and Quantum Materials, South China, aims to derive an analytical model of optical activity in black phosphorous under an external magnetic field.

Yin and his fellow authors experimented with black phosphorous—a thermodynamically stable form of phosphorus at [room temperature](#) and pressure, first synthesized in 1914—in a single, closely packed layer of atoms or a monolayer. The researchers discovered that in addition to expected strong [optical activity](#), dichroism-transmittance difference between left and right circularly polarized light—and circular birefringence, they could tune the phenomena created by altering the applied magnetic field.

The team reached their findings by deriving an analytical method to calculate the optical activity in a monolayer of black phosphorous under an external magnetic field. They were then able to obtain results showing how optical activity can be altered by changing the incident angle of the impinging light and by adjusting the magnitude of the applied magnetic field.

The results discussed in the paper show optical [activity](#) conforming to that previously observed in chiral metamaterials—material engineered to have a property that is not found in naturally occurring materials. In addition to this, they found the co-polarization transmittance increased in step with the angular frequency. The team explains the reason for this is that the conductivity of the monolayer black phosphorous decreases with increasing angular frequency, resulting in a weaker interaction with the incident light.

The researchers say that their findings could have applications in polarization optics, stereochemistry—the study of the relative spatial arrangement of atoms that form the structure of molecules and their

manipulation—and molecular biology.

More information: Chenchen Liu et al, Theory for optical activity in monolayer black phosphorus under external magnetic field, *The European Physical Journal B* (2020). [DOI: 10.1140/epjb/e2020-10390-0](https://doi.org/10.1140/epjb/e2020-10390-0)

Provided by Springer

Citation: Investigating optical activity under an external magnetic field (2020, November 6) retrieved 24 May 2024 from <https://phys.org/news/2020-11-optical-external-magnetic-field.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.