

Partial autonomy—how species separate, but not entirely

November 9 2020



Credit: AI-generated image ([disclaimer](#))

The prevailing theory until now suggests that regular gene flow through successful pairings between populations is the main barrier to their segregation into species. In recent years, however, theoretical and empirical work has shown that speciation is possible under certain circumstances despite this continued exchange of genetic information.

To date, few research results are available that show the actual extent of the phenomenon of speciation despite documented gene flow, and the time frame necessary for a new species to arise. In particular, it is not clear whether the process of speciation, once started, will inevitably lead to complete genomic and reproductive isolation sooner or later. This would mean that the diverging populations are developing into new species in the sense of the widely recognized biological species concept.

The results of a new study at the LOEWE Center for Translational Biodiversity Genomics (TBG) and the Senckenberg Biodiversity and Climate Research Center question these definitions. "We were able to show that species can achieve and maintain their dissimilarities despite the fact that they still share the majority of their genome with each other," says Dennis Schreiber, Senckenberg Biodiversity and Climate Research Center and the Johannes Gutenberg University in Mainz.

Schreiber analyzed the genome of individuals from two chironomid species, *Chironomus riparius* and *Chironomus piger*. Both species occur throughout the entire northern hemisphere and share similar ranges; however, they prefer ecologically different habitats. Their genome reveals that these species rapidly diverged several million generations (i.e., approx. 700,000 years) ago. Yet, the speciation process apparently came to a complete stop before the entire genome of each species was fully isolated from the other. Since then, hybridization, i.e., cross-breeding between individuals of both species, occurs on a regular basis.

In the process, about 70 percent of their genome is exchanged; this involves roughly half of all [genes](#). Nevertheless, the species remain ecologically recognizable, even though each can lay exclusive claim to only 30 percent of its genome (albeit including the other half of the genes), and these areas are relatively small and scattered across the entire genome. "On the one hand, this concerns genes that are related to the known ecological differences between the species, e.g., their

susceptibility to nitrogen compounds such as nitrite from liquid manure," explains Schreiber.

"At the same time, however, genes that closely cooperate with other genes on the [molecular level](#), e.g., in the respiratory chain, protein production, or transfer through membranes, also show differences. Here, incompatibility between the species would apparently be too disruptive: therefore, the species maintain their individuality in these sectors," says the head of the study, Prof. Markus Pfenninger of the LOEWE Center for Translational Biodiversity Genomics and the Senckenberg Biodiversity and Climate Research Center.

"These new findings change our concept of biological species. Apparently, this does not require an entirely independent [genome](#); instead, there are stable intermediate stages where related, well-characterized [species](#) maintain their autonomy despite a large genomic overlap. This will be subject of our future research," adds Pfenninger in conclusion.

More information: Dennis Schreiber et al. Genomic divergence landscape in recurrently hybridizing *Chironomus* sister taxa suggests stable steady state between mutual gene flow and isolation, *Evolution Letters* (2020). [DOI: 10.1002/evl3.204](https://doi.org/10.1002/evl3.204)

Provided by Senckenberg Research Institute and Natural History Museum

Citation: Partial autonomy—how species separate, but not entirely (2020, November 9) retrieved 11 May 2024 from <https://phys.org/news/2020-11-partial-autonomyhow-species.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.