

New phase of modeling the viscous coupling effects of multiphase fluid flow

November 17 2020



Predicting the multiphase permeability in pore throat by using artificial neural network. Credit: I2CNER, Kyushu University

Many applications, including carbon dioxide storage and oil recovery, involve the simultaneous flow of two or more phases of matter (solid,



liquid, gas, etc.) through porous materials. Pore-scale modeling of such multiphase flow has struggled to capture important phenomena referred to as viscous coupling effects. But now, a research team has developed a method that overcomes this limitation with potential applications to improve fuel technologies and carbon capture systems.

In a study published this month in *Advances in Water Resources*, researchers led by the International Institute for Carbon-Neutral Energy Research (WPI-I2CNER) at Kyushu University present a way to incorporate viscous coupling effects into pore-scale modeling of multiphase flow.

A common technique for studying such multiphase flows is pore network modeling (PNM), whereby simplified transport equations are solved for idealized pore geometries. PNM can be used to quickly estimate transport properties, but it neglects viscous coupling effects. An alternative approach is the lattice Boltzmann method (LBM), whereby equations governing fluid flow are solved for realistic pore geometries. Although the LBM can capture viscous coupling effects, it is extremely computationally inefficient.

The team behind this latest research had the idea to combine these two techniques. "We devised an improved model for PNM that uses data collected from LBM simulations," explains co-author of the study Takeshi Tsuji. "In the simulations, we examined multiphase flow at the pore scale for a wide range of geometric parameters and viscosity ratios."

The researchers found that for some configurations, viscous coupling effects significantly influence multiphase flow in the pore throat. They used the <u>simulation</u> results to derive a modification factor, expressed as a function of viscosity ratios, that can be easily incorporated into PNM to account for viscous coupling effects. The team also developed a machine



learning-based method to estimate the permeability associated with multiphase flow.

"We trained an artificial neural network using a database built from the results of simulations. These simulations considered different combinations of geometric parameters, viscosity ratios, and so on," says lead author Fei Jiang. "We found that the trained neural network can predict the multiphase permeability with extremely high accuracy."

This new data-driven approach not only improves PNM by including detailed <u>pore</u>-scale information, but it maintains good computational efficiency. Given that multiphase flow through porous materials is central to many natural and <u>industrial processes</u>, studies such as this one could have far-reaching implications.

More information: Fei Jiang et al, Investigation of viscous coupling effects in three-phase flow by lattice Boltzmann direct simulation and machine learning technique, *Advances in Water Resources* (2020). DOI: 10.1016/j.advwatres.2020.103797

Provided by Kyushu University, I2CNER

Citation: New phase of modeling the viscous coupling effects of multiphase fluid flow (2020, November 17) retrieved 13 May 2024 from <u>https://phys.org/news/2020-11-phase-viscous-coupling-effects-multiphase.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.