

Scientists have discovered an ancient lake bed deep beneath the Greenland ice

bed deep beneath the Greenland ice

November 10 2020, by Kevin Krajick



The largely featureless surface of the Greenland ice sheet, as seen from the window of a P3 aircraft carrying geophysical instruments aimed at detecting geologic features underneath. Credit: Kirsty Tinto/Lamont-Doherty Earth Observatory


Scientists have detected what they say are the sediments of a huge ancient lake bed sealed more than a mile under the ice of northwest Greenland—the first-ever discovery of such a sub-glacial feature anywhere in the world. Apparently formed at a time when the area was ice-free but now completely frozen in, the lake bed may be hundreds of thousands or millions of years old, and contain unique fossil and chemical traces of past climates and life. Scientists consider such data vital to understanding what the Greenland ice sheet may do in coming years as climate warms, and thus the site makes a tantalizing target for drilling. A paper describing the discovery is in press at the journal *Earth and Planetary Science Letters*.

"This could be an important repository of information, in a landscape that right now is totally concealed and inaccessible," said Guy Paxman, a postdoctoral researcher at Columbia University's Lamont-Doherty Earth Observatory and lead author of the report. "We're working to try and understand how the Greenland ice sheet has behaved in the past. It's important if we want to understand how it will behave in future decades." The ice sheet, which has been melting at an accelerating pace in recent years, contains enough water to raise global sea levels by about 24 feet.

The researchers mapped out the lake bed by analyzing data from airborne geophysical instruments that can read signals that penetrate the ice and provide images of the geologic structures below. Most of the data came from aircraft flying at low altitude over the ice sheet as part of NASA's Operation IceBridge.

The team says the [basin](#) once hosted a lake covering about 7,100 square kilometers (2,700 square miles), about the size of the U.S. states of Delaware and Rhode Island combined. Sediments in the basin, shaped vaguely like a meat cleaver, appear to range as much as 1.2 kilometers (three quarters of a mile) thick. The geophysical images show a network

of at least 18 apparent onetime stream beds carved into the adjoining bedrock in a sloping escarpment to the north that must have fed the lake. The images also show at least one apparent outlet stream to the south. The researchers calculate that the water depth in the onetime lake ranged from about 50 meters to 250 meters (a maximum of about 800 feet).




Using geophysical instruments, scientists have mapped a huge ancient lake basin (outlined here in red) below the Greenland ice, covering about 2,700 square miles). Redder colors signify higher elevations, green ones lower. A stream system incised into the bedrock that once fed the lake is shown in blue. Credit: Paxman et al., EPSL, 2020

In recent years, scientists have found existing subglacial lakes in both Greenland and Antarctica, containing liquid water sandwiched in the ice, or between bedrock and ice. This is the first time anyone has spotted a fossil lake bed, apparently formed when there was no ice, and then later covered over and frozen in place. There is no evidence that the Greenland basin contains liquid water today.

Paxman says there is no way to tell how old the lake bed is. Researchers say it is likely that ice has periodically advanced and retreated over much of Greenland for the last 10 million years, and maybe going back as far as 30 million years. A 2016 study led by Lamont-Doherty geochemist Joerg Schaefer has suggested that most of the Greenland ice may have melted for one or more extended periods some time in the last million years or so, but the details of that are sketchy. This particular area could have been repeatedly covered and uncovered, Paxman said, leaving a wide range of possibilities for the lake's history. In any case, Paxman says, the substantial depth of the sediments in the basin suggest that they must have built up during [ice-free](#) times over hundreds of thousands or millions of years.

"If we could get at those sediments, they could tell us when the ice was present or absent," he said.

The researchers assembled a detailed picture of the lake basin and its surroundings by analyzing radar, gravity and magnetic data gathered by NASA. Ice-penetrating radar provided a basic topographic map of the earth's surface underlying the ice. This revealed the outlines of the smooth, low-lying basin, nestled among higher-elevation rocks. Gravity measurements showed that the material in the basin is less dense than the surrounding hard, metamorphic rocks—evidence that it is composed of sediments washed in from the sides. Measurements of magnetism (sediments are less magnetic than solid rock) helped the team map the depths of the sediments.



A newly forming lake at the edge of the Greenland ice sheet, exposing sediments released by the ice. Such lake beds are becoming common as the ice recedes.

Credit: Kevin Krajick/Earth Institute

The researchers say the basin may have formed along a now long-dormant fault line, when the bedrock stretched out and formed a low spot. Alternatively, but less likely, previous glaciations may have carved out the depression, leaving it to fill with water when the ice receded.

What the sediments might contain is a mystery. Material washed out from the edges of the ice sheet have been found to contain the remains of pollen and other materials, suggesting that Greenland may have undergone warm periods during the last million years, allowing plants and maybe even forests to take hold. But the evidence is not conclusive, in part because it is hard to date such loose materials. The newly discovered [lake](#) bed, in contrast, could provide an intact archive of fossils and chemical signals dating to a so-far unknown distant past.

The basin "may therefore be an important site for future sub-ice drilling and the recovery of [sediment](#) records that may yield valuable insights into the glacial, climatological and environmental history" of the region, the researchers write. With the top of the sediments lying 1.8 kilometers below the current ice surface (1.1 miles), such drilling would be daunting, but not impossible. In the 1990s, researchers penetrated almost two miles into the summit of the Greenland [ice sheet](#) and recovered several feet of bedrock—at the time, the deepest ice core ever drilled. The feat, which took five years, has not since been repeated in Greenland, but a new project aimed at reaching shallower bedrock in another part of northwest Greenland is being planned for the next few years.

More information: Guy J.G. Paxman et al. A fault-bounded palaeo-lake basin preserved beneath the Greenland Ice Sheet, *Earth and Planetary Science Letters* (2020). [DOI: 10.1016/j.epsl.2020.116647](https://doi.org/10.1016/j.epsl.2020.116647)

This story is republished courtesy of Earth Institute, Columbia University

<http://blogs.ei.columbia.edu>.

Provided by Earth Institute at Columbia University

Citation: Scientists have discovered an ancient lake bed deep beneath the Greenland ice (2020, November 10) retrieved 15 May 2024 from <https://phys.org/news/2020-11-scientists-ancient-lake-bed-deep.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.