


Tunable THz radiation from 3-D topological insulator

November 2 2020, by Renae Keep




Generation of elliptically and circularly polarized terahertz beams. Credit: Haihui Zhao et al., doi 10.1117/1.AP.2.6.066003

Terahertz (THz) waves, located between the millimeter and far-infrared frequency ranges, are an electromagnetic frequency band that is as-yet incompletely recognized and understood. Xiaojun Wu of Beihang University leads a group of researchers actively seeking ways to understand, generate, and control THz radiation. Wu notes that THz waves have great potential for expanding real applications—from

imaging to information encryption—but the development of THz science and technology has been hindered by a lack of sufficiently efficient sources.

Wu's research group has been investigating a three-dimensional topological insulator of bismuth telluride (Bi_2Te_3) as a promising basis for an effective THz system. They recently systematically investigated THz radiation from Bi_2Te_3 nanofilms driven by femtosecond laser pulses. Their report published in *Advanced Photonics* demonstrates efficient generation of chiral THz waves with an arbitrarily adjustable polarization state that allows control of chirality, ellipticity, and principal axis.

According to Wu, bismuth telluride is a great candidate for future on-chip topological insulator-based terahertz systems; it has already exhibited excellent prospects in THz emission, detection, and modulation. The well-studied topological insulator presents a special spin-momentum locked surface state, which can also be accurately adjusted by various factors such as the number of atomic layers. Wu explains that this kind of THz source can efficiently radiate linearly and circularly polarized THz waves, with adjustable chirality and polarization. This will enable the development of THz science and applications in such areas as ultrafast THz opto-spintronics, polarization-based THz spectroscopy and imaging, THz biosensing, line-of-sight THz communications, and information encryption.



Schematic diagram of the polarization tunable THz emission from Bi₂Te₃. (a) Femtosecond laser pulses with horizontal linearly polarized (HLP), vertical linearly polarized (VLP), left-handed circularly polarized (LCP), and right-handed circularly polarized (RCP) illuminate onto the topological insulator Bi₂Te₃ and produce polarization tunable THz waves. (b) Macroscopic helicity-dependent photocurrents and only unidirectional spin current can be generated. (c) Microscopic electronic transition under circularly polarized laser pulse illumination. Credit: SPIE

Generation and manipulation of linearly polarized THz waves

Wu's group systematically investigated the THz radiation from topological insulator Bi₂Te₃ nanofilms driven by femtosecond laser

pulses. They found that the linearly polarized THz wave originates from the shift current formed by the ultrafast redistribution of the electron density between Bi-Te atoms in Bi_2Te_3 after the topological insulator is excited by the linearly polarized pump light. The ultrafast shift current contributes to the linearly polarized THz radiation. Due to the lattice characteristics of Bi_2Te_3 , the radiated THz waves are always linearly polarized with a three-fold rotation angle, depending on the sample azimuthal angle. This reliability makes it very convenient to arbitrarily manipulate the THz wave polarization angle by controlling the incident laser in the polarization direction.

Generation and manipulation of circularly polarized THz waves

Wu explains that, in order to produce circularly polarized THz pulses, it was necessary to simultaneously tune the pump laser polarization and the sample azimuthal angle. When the sample azimuthal angle was fixed, the researchers also obtained elliptical THz beams with various ellipticities and principle axes, due to the combination of a linear photogalvanic effect (LPGE) and a circular photogalvanic effect (CPGE), which is caused by the intrinsic time delay between the LPGE-driven and CPGE-driven THz electric field components. Within the scope of their expectations, they were able to manipulate the chirality of the emitted THz waves by varying the incident laser helicity.

Wu explains, "Helicity-dependent current is the critical reason why we can obtain spin-polarized THz pulses because we can continuously tune the magnitude and polarity of it by changing the helicity." Specific discussion of the implementation and control of circularly polarized THz radiation is included in their paper.

The authors are optimistic that their work will help further collective

understanding of femtosecond coherent control of ultrafast spin currents in light-matter interaction and will also provide an effective way to generate spin-polarized THz waves. Wu notes that the manipulation of polarization is a step toward the goal of tailoring twisted THz waves efficiently at the source.

More information: Haihui Zhao et al, Generation and manipulation of chiral terahertz waves in the three-dimensional topological insulator Bi₂Te₃, *Advanced Photonics* (2020). [DOI: 10.1117/1.AP.2.6.066003](https://doi.org/10.1117/1.AP.2.6.066003)

Provided by SPIE

Citation: Tunable THz radiation from 3-D topological insulator (2020, November 2) retrieved 3 May 2024 from <https://phys.org/news/2020-11-tunable-thz-d-topological-insulator.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--