


Cell membranes in super resolution

December 2 2020



Sphingolipid expansion microscopy (ExM) of tenfold expanded cells infected with chlamydia. The bacterial membranes are marked green; the inner and outer membranes of the bacteria can be distinguished (c). Under (a) confocal laser scanning and under (b) structured illumination microscopy (SIM). Scale bars: 10 and 2 microns in the small white rectangles respectively. Credit: Sauer group / University of Würzburg

Expansion microscopy (ExM) enables the imaging of cells and their components with a spatial resolution far below 200 nanometres. For this purpose, the proteins of the sample under investigation are cross-linked into a swellable polymer. Once the interactions between the molecules

have been destroyed, the samples can be expanded many times over with water. This allows detailed insights into their structures.

"This method was previously limited to proteins. In the journal *Nature Communications* we are now presenting a way of expanding lipids and thus cell membranes," says Professor Markus Sauer, an expert in super-resolution microscopy from the Biocentre of Julius-Maximilians-Universität (JMU) Würzburg in Bavaria, Germany. JMU professors Thomas Rudel (microbiology) and Jürgen Seibel (chemistry) are also involved in the publication.

Synthetic lipids are marked and expanded

Jürgen Seibel's team has synthesized functionalised sphingolipids, which are an important component of cell membranes. If these lipids are added to [cell cultures](#), they are incorporated into the cell membranes. They can then be marked with a dye and expanded four to ten times in a swellable polymer.

The JMU researchers show that this method—in combination with structured illumination microscopy (SIM) – makes it possible for the first time to image different membranes and their interactions with proteins with a resolution of 10 to 20 nanometres: cell membranes, the outer and inner cell nuclear membrane and also the membranes of intracellular organelles such as mitochondria.

Focusing on bacteria and viruses

The sphingolipids also integrate highly efficiently into the membranes of bacteria. This means that, for the first time, pathogens such as *Neisseria gonorrhoeae*, *Chlamydia trachomatis* and *Simkania negevensis* can now be visualized in infected [cells](#) with a resolution that was previously only

achieved using electron microscopy. Even the inner and outer membranes of Gram-negative bacteria can be distinguished from each other.

"With the new super-resolution microscopic methods, we now want to investigate bacterial infection mechanisms and causes of antibiotic resistance. What we learn in the process could possibly be used for improved therapies," says Professor Thomas Rudel, an expert on bacterial infections.

The sphingolipids might also integrate into the [membrane](#) of viruses. If this is successful, the interactions of corona viruses with cells could be studied for the first time with high resolution light microscopy.

More information: Ralph Götz et al. Nanoscale imaging of bacterial infections by sphingolipid expansion microscopy, *Nature Communications* (2020). [DOI: 10.1038/s41467-020-19897-1](https://doi.org/10.1038/s41467-020-19897-1)

Provided by Julius-Maximilians-Universität Würzburg

Citation: Cell membranes in super resolution (2020, December 2) retrieved 2 May 2024 from <https://phys.org/news/2020-12-cell-membranes-super-resolution.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--