

A groggy climate giant: subsea permafrost is still waking up after 12,000 years

December 22 2020



The coastline of the Bykovsky Peninsula in the central Laptev Sea, Siberia retreats during summer, when ice-rich blocks of permafrost fall to the beach and are eroded by waves. Credit: 2017, P. Overduin

In the far north, the swelling Arctic Ocean inundated vast swaths of coastal tundra and steppe ecosystems. Though the ocean water was only a few degrees above freezing, it started to thaw the permafrost beneath it, exposing billions of tons of organic matter to microbial breakdown.

The decomposing organic matter began producing CO₂ and CH₄, two of the most important greenhouse gases.

Though researchers have been studying degrading subsea permafrost for decades, difficulty collecting measurements and sharing data across international and disciplinary divides have prevented an overall estimate of the amount of [carbon](#) and the rate of release. A new study, led by Ph.D. candidate Sara Sayedi and senior researcher Dr. Ben Abbott at Brigham Young University (BYU) published in IOP Publishing journal *Environmental Research Letters*, sheds light on the subsea permafrost climate feedback, generating the first estimates of circumarctic carbon stocks, greenhouse gas release, and possible future response of the subsea permafrost zone.

Sayedi and an international team of 25 permafrost researchers worked under the coordination of the Permafrost Carbon Network (PCN), which is supported by the U.S. National Science Foundation. The researchers combined findings from published and unpublished studies to estimate the size of the past and present subsea carbon stock and how much greenhouse gas it might produce over the next three centuries.


Using a methodology called expert assessment, which combines multiple, independent plausible values, the researchers estimated that the subsea permafrost region currently traps 60 billion tons of methane and contains 560 billion tons of organic carbon in sediment and soil. For reference, humans have released a total of about 500 billion tons of carbon into the atmosphere since the Industrial Revolution. This makes the subsea permafrost carbon stock a potential giant ecosystem feedback to climate change.

"Subsea permafrost is really unique because it is still responding to a dramatic climate transition from more than ten thousand years ago," Sayedi said. "In some ways, it can give us a peek into the possible

response of permafrost that is thawing today because of [human activity](#)."

Estimates from Sayedi's team suggest that subsea permafrost is already releasing substantial amounts of greenhouse gas. However, this release is mainly due to ancient climate change rather than current human activity. They estimate that subsea permafrost releases approximately 140 million tons of CO₂ and 5.3 million tons of CH₄ to the atmosphere each year. This is similar in magnitude to the overall greenhouse gas footprint of Spain.

The researchers found that if human-caused [climate change](#) continues, the release of CH₄ and CO₂ from subsea permafrost could increase substantially. However, this response is expected to occur over the next three centuries rather than abruptly. Researchers estimated that the amount of future greenhouse gas release from subsea permafrost depends directly on future human emissions. They found that under a business-as-usual scenario, warming subsea permafrost releases four times more additional CO₂ and CH₄ compared to when human emissions are reduced to keep warming less than 2°C.




Artistic diagram of the subsea and coastal permafrost ecosystems, emphasizing greenhouse gas production and release. Credit: Original artwork created for this study by Victor Oleg Leshyk at Northern Arizona University.

"These results are important because they indicate a substantial but slow climate feedback," Sayedi explained. "Some coverage of this region has

suggested that human emissions could trigger catastrophic release of methane hydrates, but our study suggests a gradual increase over many decades."

Even if this climate feedback is relatively gradual, the researchers point out that subsea permafrost is not included in any current climate agreements or greenhouse gas targets. Sayedi emphasized that there is still a large amount of uncertainty about subsea permafrost and that additional research is needed.



The coastline of the Bykovsky Peninsula in the central Laptev Sea, Siberia retreats during summer, when ice-rich blocks of permafrost fall to the beach and are eroded by waves. Credit: 2017, P. Overduin

"Compared to how important subsea [permafrost](#) could be for future [climate](#), we know shockingly little about this ecosystem," Sayedi said.

"We need more sediment and soil samples, as well as a better monitoring network to detect when [greenhouse](#) gas release responds to current warming and just how quickly this giant pool of carbon will wake from its frozen slumber."

More information: Sayedeh Sara Sayedi et al, Subsea permafrost carbon stocks and climate change sensitivity estimated by expert assessment, *Environmental Research Letters* (2020). [DOI: 10.1088/1748-9326/abcc29](#)

Provided by Institute of Physics

Citation: A groggy climate giant: subsea permafrost is still waking up after 12,000 years (2020, December 22) retrieved 11 July 2024 from <https://phys.org/news/2020-12-groggy-climate-giant-subsea-permafrost.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.
