


Near-atomic-scale analysis of frozen water

December 11 2020, by Thamarasee Jeewandara




Summary of the atom probe data from a thick layer of ice. (A) Mass spectrum of acquired APT dataset of D₂O ice at 100 pJ, 200 kHz, and a detection rate of 0.5%. (B) Sectioned mass spectrum from (A) to illustrate D_xH_{3-x}O complex peaks. (C) 3D reconstruction map of D₂O. Inset capture shows SEM image of the specimen. Credit: Science Advances, doi: 10.1126/sciadv.abd6324

Advances in transmission electron microscopy (TEM) can allow [cryo-imaging](#) of biological and biochemical systems in liquid form, however, such approaches do not possess advanced analytical capabilities. In a new report now published on *Science Advances*, A. A. El-Zoka and an international team of researchers in Germany, Canada, France, and the U.K., used [atom probe tomography](#) to analyze frozen liquids in three-dimensions (3-D) with sub-nanometer scale resolution. In this work, the team first introduced a specimen preparation strategy using nano-porous

gold and used ice formed from high-purity [deuterated water](#) (hard water) alongside a solution of sodium chloride (50 mM) dissolved in high-purity deuterated water. They then analyzed the gold-ice interface to reveal increased solute concentrations across the interface. The scientists explored a range of experimental conditions to understand atom probe analyses of bulk aqueous specimens. Then they discussed the physical processes associated with the observed phenomena. The study showed the practicality of using frozen water as a carrier for near-atomic-scale analyses of objects in solution via atom probe tomography.

Transmission electron microscopy and atom probe tomography

Transmission electron microscopy (TEM) has undergone significant progress in recent decades, partly leading to the [2017 Nobel prize in Chemistry](#), due to the innovation of [cryo-electron microscopy](#) (cryo-EM) to determine the high-resolution structure of biomolecules in solution. The cryo-EM technique notably offers the ability to freeze specimens rapidly so that water molecules present in the specimens turn into transparent ice crystals. Tremendous parallel efforts have similarly established atomically resolved electron tomography methods to accomplish groundbreaking discoveries [in materials science](#). Despite the powerful analytical capabilities, the approaches cannot readily measure the atomic-scale composition of a specimen. Here, El-Zoka et al. described the analysis of micron-thick layers of frozen water formed on nanoporous gold (NPG), with typical applications in catalysis, electrochemical sensing and actuation due to [a high surface-area-to-volume ratio](#) and gold-rich surface. The team therefore used NPG as a hydrophilic (water-loving) substrate on which to analyze ice using atom [probe](#) tomography.



SEM images of in situ APT specimen preparation of an ice sample on NPG (nanoporous gold). (A) The 200- and 75- μm ion beam annular patterns for outer and inner diameters, respectively, were made on the ice/NPG sample. (B) The ice/NPG pillar was milled until the height of the Au post reached

Citation: Near-atomic-scale analysis of frozen water (2020, December 11) retrieved 26 April 2024 from <https://phys.org/news/2020-12-near-atomic-scale-analysis-frozen.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.