


A neglected mechanism in antiferromagnets may be key to spintronics

December 7 2020



Optical laser pulses generate two types of torque, field-like (τ_{FL}) and damping-like (τ_{DL}), that affect the three magnetizations of $YMnO_3$. The damping-like torque has a pronounced effect on the inherent elliptically oscillating magnetizations, producing a large instantaneous change in antiferromagnetic ordering. Credit: Tokyo Tech

Enormous efforts are being made worldwide in a technological field that

could far surpass the capabilities of conventional electronics: Spintronics. Instead of operating based on the collective movement of charged particles (electrons), spintronic devices could perform memory storage and data transmission by manipulating spin, an intrinsic property of elementary particles related to angular momentum and from which many magnetic characteristics in materials arise. Unfortunately, controlling spin has proven to be a challenging endeavor, leading physicists and engineers to look for efficient materials and techniques to do so.

In this regard, antiferromagnetic materials (AFMs) are good candidates for spintronics because they are resistant to external magnetic fields and allow for switching spin values in timescales of picoseconds. One promising strategy to manipulate spin orientation in AFMs is using an optical laser to create extremely short-lived magnetic field pulses, a phenomenon known as the inverse Faraday effect (IFE). Although the IFE in AFMs generates two very distinct types of [torque](#) (rotational force) on their magnetization, it now seems the most important of the two has somehow been neglected in research.

In a recent study published in *Nature Communications*, a trio of scientists, including Professor Takuya Satoh from the Tokyo Tech, Japan, delved deep into this issue. Spin dynamics in AFMs are described by a sum of two terms: field-like torque and damping-like torque. The latter, as the word 'damping' implies, is related to the gradual decay (or dying off) of the spin oscillations triggered by the optical pulses on the material.

Until now, scientists studied the damping-like torque only from the perspective of spin relaxation after excitation, believing its amplitude to be small during the ultrashort spin excitation process. In this study, however, Prof Satoh and colleagues found it to be, in some cases, the main player in terms of spin reorientation due to the IFE. Through

theoretical analyses and experimental verification in both YMnO₃ and HoMnO₃, they clarified the conditions under which the damping effect becomes the dominant spin excitation mechanism.

A simplified interpretation of the findings can be as follows. Imagine a hanging pendulum (magnetization direction) oscillating in wide arcs, drawing a very pronounced ellipse. The damping-like torque produces a large instantaneous perturbation in the direction of the small diameter, 'tipping it off' and causing it to lean like a spinning top that is about to fall. "The otherwise small damping-related magnetization causes a large spin canting because of the extreme ellipticity inherent to AFMs," explains Prof Satoh. "Considering that it is possible to adjust the strength of the damping by strategically selecting the ions in the AFM, we might have found a way to tune material properties for specific spintronic applications," he adds.

The trio of scientists also tested how spin dynamics are influenced by temperature, which affects and even destroys antiferromagnetic order past certain thresholds. By putting the materials close to the critical transition points, they managed to produce a more pronounced effect from damping-type torque. Excited about the results, Prof Satoh remarks: "Our results indicate that optically generated torques might provide the long sought-after tool enabling the efficient realization of ultrafast spin switching in AFMs."

Although much more research will certainly be needed before applied spintronics becomes a reality, uncovering efficient mechanisms for [spin](#) manipulation is obviously among the first steps. This study proves that such mechanisms might be hidden in phenomena we know and neglect!

More information: Christian Tzschaschel et al, Efficient spin excitation via ultrafast damping-like torques in antiferromagnets, *Nature Communications* (2020). [DOI: 10.1038/s41467-020-19749-y](https://doi.org/10.1038/s41467-020-19749-y)

Provided by Tokyo Institute of Technology

Citation: A neglected mechanism in antiferromagnets may be key to spintronics (2020, December 7) retrieved 6 May 2024 from <https://phys.org/news/2020-12-neglected-mechanism-antiferromagnets-key-spintronics.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.