


# Scientists reveal reaction mechanism of $^{11}\text{Be}$ nucleus

December 8 2020, by Zhang Nannan



Experimental differential breakup cross section for  $^{11}\text{Be}+^{208}\text{Pb}$  system at  $E_{\text{lab}}=140$  MeV compared with the CDCC and XCDCC calculations. Credit: DUAN Fangfang

Scientists from the Institute of Modern Physics (IMP) of the Chinese Academy of Sciences (CAS) and their collaborators have lately made new progress in the study of the reaction mechanism of  $^{11}\text{Be}$  nucleus. The study will help understand the effect of exotic structures such as the neutron halo on the reaction characteristics.

Elastic and breakup reactions induced by stable as well as unstable atomic nuclei constitute a fruitful area of research in [nuclear physics](#). The angular distributions of elastic scattering cross sections can exhibit different features depending on the incident [energy](#) and the structure of the colliding nuclei.

$^{11}\text{Be}$  is an archetype of one-neutron halo nucleus, whose valence neutron has a low binding energy. In previous studies, measurements of the elastic scattering of  $^{11}\text{Be}$  have been made at energies around the Coulomb barrier on several medium to heavy targets.

For the first time, researchers at IMP and their collaborators have studied the elastic scattering and breakup reactions of  $^{11}\text{Be}$  on  $^{208}\text{Pb}$  at an incident energy of 140 MeV, which corresponds to about 3.5 times of the Coulomb barrier ( $V_B \approx 39.5$  MeV). The experiment was conducted at the Radioactive Ion Beam Line in Lanzhou, the Heavy Ion Research Facility of Lanzhou (HIRFL-RIBLL).

A strong suppression of the Coulomb nuclear interference peak is observed in the measured quasielastic scattering angular distribution. It is the first time that researchers have demonstrated the persistence of the strong breakup coupling effect for reaction systems involving neutron-halo nuclei at this relatively high incident energy.

By calculations, they predicted that the effect of the core excitation of the reaction system can be ignored.

Moreover, the angular and energy distributions of the  $^{10}\text{Be}$  fragments have been measured for the first [time](#). These calculations revealed that the fragments are mostly produced by an elastic breakup (EBU) mechanism. However, the contribution of nonelastic breakup (NEB) is also found in the data. The results also reveal that these fragments experience a post-acceleration effect after the [breakup](#) takes place.

Provided by Chinese Academy of Sciences

Citation: Scientists reveal reaction mechanism of  $^{11}\text{Be}$  nucleus (2020, December 8) retrieved 11 May 2024 from <https://phys.org/news/2020-12-scientists-reveal-reaction-mechanism-11be.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.