

# Synergy between biotech and classical control tactics rid US of invasive pest

December 21 2020

---



Releasing infertile adult pink bollworm moths from airplanes by the billions to disrupt reproduction of moths in the wild was an important factor in eradicating this pest in the U.S. in 2018. Credit: Agricultural Research Service-USDA.

Genetically engineered cotton and classical pest control tactics combined to rid the United States and Northern Mexico of a devastating pest, according to a new study by Agricultural Research Service (ARS) and

University of Arizona (UofA) scientists published in the *Proceedings of the National Academy of Sciences*.

For most of the past century, the [pink bollworm](#) was the major [cotton](#) pest in the Southwest. For decades, cotton growing in Arizona, California, Texas, and New Mexico was only possible because farmers sprayed pesticides an average of 12 times a year, nine specifically against pink bollworm. Some farmers sprayed as often as 25 times a year without reaching control. In 1990, pink bollworm cost cotton growers \$48 million in Arizona alone.

A coordinated and multitactical list of areawide and integrated pest management strategies were developed over the years in hopes of putting down this pest while replacing expensive and environmentally hazardous chemical pesticides, explained research entomologist Jeffrey Fabrick, one of the authors of the study. Fabrick is with the ARS-USDA Pest Management and Biocontrol Research Unit in Maricopa, Arizona.

"By analyzing computer simulations and 21 years of field data from Arizona, we proved that genetically engineered cotton and release of billions of sterile pink bollworm moths acted synergistically to suppress this pest," Fabrick said.

Both the [computer simulations](#) and what was seen in the field from 2006 to 2010 showed neither of the two tactics would have worked if used alone, he added.



Pink bollworm caterpillars emerging from a damaged cotton boll. Credit: Agricultural Research Service-USDA

"Collaboration among farmers and scientists from government, industry, and academia was essential for the remarkable success of the pink bollworm eradication program," said Bruce Tabashnik, lead author of the study and regents professor in the UofA Department of Entomology.

In the late 1960s and 1970s, ARS scientists first began powering up the fight against pink bollworm. They helped create the artificial pheromones that allowed precise tracking of the pest as well as the first synthetic diet and methods for raising sterile pink bollworm moths to disrupt mating. Releasing synthetic female sex pheromone in [cotton fields](#) also was used to confuse males and disrupt mating. Another important tactic required farmers to plow down cotton residues after

harvest to reduce overwintering survival of pink bollworm.

Enter genetically modified Bt cotton in 1996. Bt cotton is engineered to produce one or more proteins from the bacterium *Bacillus thuringiensis* (Bt for short) that kills pink bollworm and other related caterpillar pests and are harmless to people and most other insects, unlike broad spectrum pesticides. Growing mostly Bt cotton knocked the pink bollworm population down by 90 percent in 10 years. At the same time, farmers continued employing other techniques.

By 2006, for the first time, eradication became a practical reality. With an eye to finishing off pink bollworm, detailed cooperative plans were developed by a coalition that included cotton farmers, grower organizations, ARS researchers, USDA's Animal and Plant Health Inspection Service (APHIS), the biotech industry, the Arizona Department of Agriculture, the Arizona Cotton Research and Protection Council, and UofA extension and research scientists. Many of these groups' counterparts in Northern Mexico were also recruited.

APHIS also scaled up production of sterile pink bollworm moths so that billions of them were unleashed by airplanes to overwhelm any field populations of the pest.

Removal of pink bollworm saved U.S. cotton farmers \$192 million from 2014 to 2019 alone, according to the study. Pink bollworm suppression has also facilitated integrated [pest](#) management for all other cotton pests. Overall, this reduced [insecticide use](#) by 82 percent, avoiding application of 25 million pounds of insecticides in Arizona alone during the past two decades. It improved the overall environment and brought back beneficial insects as the ecology returned to a more natural balance.

**More information:** Bruce E. Tabashnik et al., "Transgenic cotton and sterile insect releases synergize eradication of pink bollworm a century

after it invaded the United States," *PNAS* (2020).  
[www.pnas.org/cgi/doi/10.1073/pnas.2019115118](http://www.pnas.org/cgi/doi/10.1073/pnas.2019115118)

Provided by United States Department of Agriculture

Citation: Synergy between biotech and classical control tactics rid US of invasive pest (2020, December 21) retrieved 24 May 2024 from <https://phys.org/news/2020-12-synergy-biotech-classical-tactics-invasive.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.