


# Spilling the beans on coffee's true identity

January 13 2021

---



Credit: Pixabay/CC0 Public Domain

People worldwide want their coffee to be both satisfying and reasonably priced. To meet these standards, roasters typically use a blend of two types of beans, arabica and robusta. But, some use more of the cheaper robusta than they acknowledge, as the bean composition is difficult to determine after roasting. Now, researchers reporting in *ACS' Journal of*

*Agricultural and Food Chemistry* have developed a new way to assess exactly what's in that cup of joe.

Coffee blends can have good quality and flavor. However, arabica beans are more desirable than other types, resulting in a higher market value for blends containing a higher proportion of this variety. In some cases, producers dilute their blends with the less expensive robusta beans, yet that is hard for consumers to discern. Recently, methods involving chromatography or spectroscopy were developed for [coffee](#) authentication, but most of these are labor- and time-intensive, or use chloroform for the extraction, which limits the types of [compounds](#) that can be detected. In some studies, researchers used [nuclear magnetic resonance](#) (NMR) spectroscopy to monitor the amount of 16-*O*-methylcafestol (16-OMC) in coffee, but its concentrations vary depending on geographic location and cultivar. So, Fabrice Berru  and colleagues wanted to build on their previous work with NMR to assess the chemical make-up of each coffee bean variety and confirm the blends of real samples.

The researchers extracted compounds from a test set of pure coffee and known blends with methanol and identified the compounds with NMR. The team found 12 compounds with measurable concentrations, and two had significantly different amounts between the coffee varieties. Elevated concentrations of 16-OMC were unique to robusta, while high concentrations of kahewol—a compound previously found in coffee beans by other researchers—were distinct in arabica. There was a direct, reproducible relationship between 16-OMC and kahewol concentrations found in the blends of the two varieties. The team then measured 16-OMC and kahewol levels, in addition to other flavor molecules, in 292 samples from producers around the world. They could successfully authenticate pure coffee, even with relatively low concentrations of the two indicator compounds. For samples in which the composition of blends was known, the team's predictions were within 15% of the actual

ratio. The new method results in a more robust and reliable way to verify unadulterated coffee and predict blends than previously reported approaches, the researchers say.

**More information:** Ian W. Burton et al. Quantitative NMR Methodology for the Authentication of Roasted Coffee and Prediction of Blends, *Journal of Agricultural and Food Chemistry* (2020). [DOI: 10.1021/acs.jafc.0c06239](https://doi.org/10.1021/acs.jafc.0c06239)

Provided by American Chemical Society

Citation: Spilling the beans on coffee's true identity (2021, January 13) retrieved 22 June 2024 from <https://phys.org/news/2021-01-beans-coffee-true-identity.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.