


Catalyst transforms plastic waste to valuable ingredients at low temperature

January 5 2021



Credit: Osaka City University

For the first time, researchers have used a novel catalyst process to recycle a type of plastic found in everything from grocery bags and food packaging to toys and electronics into liquid fuels and wax.

The team published their results on Dec. 10 in *Applied Catalysis B: Environmental*.

"Plastics are essential materials for our life because they bring safety and hygiene to our society," said paper co-authors Masazumi Tamura, associate professor in the Research Center for Artificial Photosynthesis

in the Advanced Research Institute for Natural Science and Technology in Osaka City University, and Keiichi Tomishige, professor in the Graduate School of Engineering in Tohoku University. "However, the growth of the global [plastic](#) production and the rapid penetration of plastics into our society brought mismanagement of waste plastics, causing serious environmental and biological issues such as ocean pollution."

Polyolefinic plastics—the most common plastic—have [physical properties](#) that make it difficult for a catalyst, responsible for inducing chemical transformation, to interact directly with the molecular elements to cause a change. Current recycling efforts require temperatures of at least 573 degrees Kelvin, and up to 1,173 degrees Kelvin. For comparison, water boils at 373.15 degrees Kelvin, and the surface of the Sun is 5,778 degrees Kelvin.

The researchers looked to heterogenous catalysts in an effort to find a reaction that might require a lower temperature to activate. By using a catalyst in a different state of matter than the plastics, they hypothesized that the reaction would be stronger at a lower temperature.

They combined ruthenium, a metal in the platinum family, with cerium dioxide, used to polish glass among other applications, to produce a catalyst that caused the plastics to react at 473 degrees Kelvin. While still high for human sensibilities, it requires significantly less energy input compared to other catalyst systems.

According to Tamura and Tomishige, ruthenium-based catalysts have never been reported in the [scientific literature](#) as a way to directly recycle polyolefinic plastics.

"Our approach acted as an effective and reusable heterogeneous catalyst, showing much higher activity than other metal-supported catalysts,

working even under mild reaction conditions," Tamura and Tomishige said. "Furthermore, a [plastic bag](#) and waste plastics could be transformed to valuable chemicals in [high yields](#)."

The researchers processed a plastic bag and [waste](#) plastics with the catalyst, producing a 92% yield of useful materials, including a 77% yield of liquid fuel and a 15% yield of wax.

"This [catalyst](#) system is expected to contribute to not only suppression of plastic wastes but also to utilization of plastic wastes as raw materials for production of chemicals," Tamura and Tomishige said.

More information: Yosuke Nakaji et al. Low-temperature catalytic upgrading of waste polyolefinic plastics into liquid fuels and waxes, *Applied Catalysis B: Environmental* (2020). [DOI: 10.1016/j.apcatb.2020.119805](#)

Provided by Osaka City University

Citation: Catalyst transforms plastic waste to valuable ingredients at low temperature (2021, January 5) retrieved 10 July 2024 from <https://phys.org/news/2021-01-catalyst-plastic-valuable-ingredients-temperature.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
