

## Drone-based early detection system to monitor changes in the Amazon

January 6 2021



A drone flies toward a rainstorm in the Amazon. Credit: Jianhuai Ye/ Harvard SEAS

The Amazon basin, home to the largest rainforest in the world, plays a crucial role in maintaining the planet's carbon budget, absorbing and storing billions of tons of carbon dioxide annually. But a tipping point looms—one that may turn this vital carbon sink into one of the largest sources of carbon dioxide on the planet.



By "smelling the forest," a Harvard-led team of researchers funded by the U.S. National Science Foundation is attempting to measure how and when that change could happen. The scientists report their results in the journal *Environmental Science: Atmospheres*.

As the planet warms, drought, wildfire and changing weather patterns threaten some 400 billion trees in the Amazon, some of which are already at risk from logging and mining. As trees are damaged or killed, they decompose and release carbon into the atmosphere.

"Climate change, as well as human-caused deforestation and biomass burning, can lead to ecological and climatic tipping points that could release massive pools of stored <u>carbon</u>," said atmospheric scientist Scot Martin of Harvard University.

The question is: How do we know when we're approaching that tipping point?

Martin, with an international team of researchers and collaborators from Amazonas State University and the Amazonas State Research Support Foundation, has developed an early detection system to monitor changes in the Amazon.

The research aims to better understand how the Amazon responds to stress. The team is mapping and monitoring the unique chemical signals emitted by trees known as volatile organic compounds, or VOCs.

Every species of plant emits a different VOC signature, like a fingerprint, which can change based on the season or stresses to the plant, for example, drought or floods.

"Forests can speak to us through VOCs," said Martin. "Translating these signals can lead to an understanding of how forest ecosystems respond to



climate stress and climate change."

But there are major challenges to collecting data on VOCs. Airplanes can cover large distances but can't fly low enough to collect VOC samples, which reach heights of only a kilometer or less above the canopy. Towers can sense at the right height but only for the local ecosystem.

To bridge this canyon of data, the team is turning to drones.

"What makes drone-based sensors so exciting is that they offer the possibility of collecting data at unexplored scales," said Martin. "This could lead to revolutionary insights into Amazonian ecosystems under climate stress and anticipated changes in climate and biodiversity."

Added Sylvia Edgerton, a program director in NSF's Division of Atmospheric and Geospace Sciences, "Using an unmanned aerial vehicle to map concentrations of these compounds throughout the canopy is a novel approach for assessing how much of their concentration comes from <u>biological processes</u> in plants versus secondary production from atmospheric chemical reactions in the canopy."

**More information:** Jianhuai Ye et al. Near-canopy horizontal concentration heterogeneity of semivolatile oxygenated organic compounds and implications for 2-methyltetrols primary emissions, *Environmental Science: Atmospheres* (2020). DOI: 10.1039/D0EA00006J

## Provided by National Science Foundation

Citation: Drone-based early detection system to monitor changes in the Amazon (2021, January 6) retrieved 25 June 2024 from <u>https://phys.org/news/2021-01-drone-based-early-amazon.html</u>



This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.