


Going with the grains to explain a fundamental tectonic force

January 19 2021, by Jim Shelton



Mylonite is a fine-grained, compact metamorphic rock produced by dynamic recrystallization of the constituent minerals resulting in a reduction of the grain size of the rock. Credit: Wikipedia

A new study suggests that tiny, mineral grains—squeezed and mixed over millions of years—set in motion the chain of events that plunge massive tectonic plates deep into the Earth's interior.

The theory, proposed by Yale scientists David Bercovici and Elvira Mulyukova, may provide an [origin story](#) for [subduction](#), one of the most [fundamental forces](#) responsible for the dynamic nature of the planet.


The study appears in the *Proceedings of the National Academy of Sciences*.

Subduction occurs when one tectonic plate slides underneath another plate and then sinks into the Earth's mantle. Its role in major geological processes is immense: It is the main engine for tectonic motion. It builds mountains, triggers earthquakes, forms volcanoes, and drives the geologic carbon cycle.

Yet researchers have been uncertain about what initiates subduction.

"Why Earth even has subduction, unlike other [terrestrial planets](#) as far as we know, is a mystery," said Bercovici, Yale's Frederick William Beinecke Professor and chair of Earth and Planetary Sciences.

"Mantle rock near the surface that has cooled for hundreds of millions of years has two competing effects," he said. "While it's gotten colder and heavier and wants to sink, it's also gotten stiffer and doesn't want to sink. The stiffening effect should win out, as it does on most planets, but on Earth, for some reason, it doesn't."



A conceptual sketch of the ocean basin setting for the new model. Inset images from a computer model show mineral fraction, grain size, and weakness. Credit: Elvira Mulyukova and David Bercovici

According to the theoretical model developed by Bercovici and Mulyukova, a research scientist at Yale, subduction may initiate at the margins between Earth's sea floor and continents.

The model shows that tectonic stresses in an oceanic plate cause its [mineral grains](#) to mix with each other, become damaged, and eventually shrink. Over a period of approximately 100 million years, this process weakens the oceanic plate and makes it susceptible to vertical shear and bending—which are associated with the start of subduction.

"The real bottleneck for tectonic plate activity on a terrestrial planet is how fast its massive, rocky layers can deform," said Mulyukova. "The

rocks can deform only as fast as their tiny mineral grains allow. Our model explains how these changes in mineral grains can dramatically weaken the rock and make subduction possible on a planet like Earth."

More information: David Bercovici et al., "Evolution and demise of passive margins through grain mixing and damage," *PNAS* (2021).
www.pnas.org/cgi/doi/10.1073/pnas.2011247118

Provided by Yale University

Citation: Going with the grains to explain a fundamental tectonic force (2021, January 19)
retrieved 25 April 2024 from <https://phys.org/news/2021-01-grains-fundamental-tectonic.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.