

Increasing soil pH reduces fertilizer-derived nitrous oxide emissions

January 29 2021



Experimental plots with greenhouse gas measurement chambers in Teagasc Johnstown Castle Research Centre. Credit: Teagasc

A new scientific paper from Teagasc has shown that getting soil pH right through a liming program can significantly reduce emissions of nitrous oxide (N_2O), a potent greenhouse gas. The paper from researchers at the Teagasc Environment, Soils and Land-Use Department in Johnstown Castle, County Wexford has just been published in the scientific journal

Agriculture, Ecosystems and Environment (AGEE). It concludes that increasing soil pH reduces fertilizer derived N₂O emissions in intensively managed temperate grassland.

Head of the Teagasc Environment Research Department, Dr. Karl Richards said: "The paper shows that there are reduced N₂O emissions from fertilizer applied to higher pH soils, where the pH is in the recommended agronomic range. Farmers that can improve soil pH for agronomic benefits, can also reduce N₂O emissions. This represents a win-win for the farmer and the environment."

Soil pH is generally considered a master variable, controlling a wide range of physical, chemical and biological properties, including a significant effect on microbial processes responsible for production and consumption of N₂O. Senior Research officer at Johnstown Castle, Dr. David Wall stated that "using an existing long-term intensive [grassland](#) liming and P trial, this research investigated the effect of longer-term lime and P management and their interaction on N₂O emissions and grassland productivity.

Postdoc Researcher with Teagasc, Ognjen (Oggy) Zurovec outlined the main findings: "We found that a long-term increase in soil pH as a result of liming significantly decreased N₂O emissions over 12-month measurement period. In addition, keeping the soil pH and P at the optimum level has the potential to further reduce N₂O emissions due to higher grass N uptake through increased yields. This means that the application of good farming practices has considerable N₂O mitigation potential in temperate grasslands."

The results showed that applying 5 tons of lime per hectare every three to four years increased soil pH from 5.1 to 6.9 and reduced N₂O emissions by 39%. The study estimated that the increase in [soil](#) pH of grasslands in Ireland over the last 12 years potentially reduced national

N₂O emissions by 95,000 T CO₂-eq yr⁻¹, with potential for a further reduction by up to 254,000 T CO₂-eq yr⁻¹ if all the remaining acidic soils are brought up to optimal pH.

More information: Ognjen Žurovec et al. Increasing soil pH reduces fertiliser derived N₂O emissions in intensively managed temperate grassland, *Agriculture, Ecosystems & Environment* (2021). [DOI: 10.1016/j.agee.2021.107319](https://doi.org/10.1016/j.agee.2021.107319)

Provided by Teagasc

Citation: Increasing soil pH reduces fertilizer-derived nitrous oxide emissions (2021, January 29) retrieved 19 April 2024 from <https://phys.org/news/2021-01-soil-ph-fertilizer-derived-nitrous-oxide.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.