

## Ultraviolet light helps marine animals to tell the time of year

January 11 2021



Post-doc and co-author Enrique Arboleda retrieving light measurement data. Credit: © Enrique Arboleda/Markus Hartl

Most organisms on Earth depend on the energy from the sun. Sunlight is also an important coordinator of life's timers. Animals take important cues for proliferation, activity, feeding, or sleep from changing light conditions. These rhythms also exist in humans—as changing light conditions across the year can strongly impact human mood and



## psychology.

Part of the natural light from the sun we are exposed to consists of ultraviolet (UVA and UVB) light, a short-wavelength part of the spectrum that is largely missing in artificial lighting. So far, most research on seasonal cycles has focused on daylength. "In contrast to previous assumptions, we discovered that, in addition to daylength, the intensity of UVA light influences the seasonal responses of the bristle worm Platynereis dumerilii," explains first author Vinoth Babu Veedin Rajan, who carried out the work during his Ph.D. in the Tessmar-Raible lab.

Initially, the researchers set out to measure the <u>light conditions</u> the <u>worms</u> are exposed to in the Bay of Naples (Italy), one of their <u>natural habitats</u>. Upon inspection of the data the scientists noticed that the light intensities of UV wavelengths exhibit annual changes distinct from the variation in daylength. They wondered if these variations in <u>light</u> <u>intensity</u> could be used by the worms to tell the time of the year. In cooperation with an <u>electrical engineer</u>, the team built lamps that closely resemble the natural light environment, including UVA. The intensity of UVA light altered worm behavior. The scientists then identified a key light receptor (c-opsin 1) involved in the response. Worms lacking the receptors showed altered levels of enzymes that produce important neurohormones such as dopamine and serotonin.

UVA-sensitive opsins with similar properties are also found in other organisms, ranging from fish to humans. "We like to think of UVA light as something that we should be protected from, but in these <u>marine</u> <u>worms</u> it clearly has a biological function. It will be interesting to find out what impact UVA levels have on other animals, and even ourselves," says Tessmar-Raible.

As part of a complex environment, the correct timing of the behavior of



marine life also has an impact on populations and the ecosystem of the ocean. How UVA light exactly influences this delicate balance is still unclear. "Understanding more about the relevance of UVA <u>light</u> for marine animal physiology and behavior will therefore also be of ecological importance," the researchers suggest. This latter aspect has already prompted interest from other marine chrono-ecologists, which may pave the way for future collaborations to bridge molecular genetics to ecology.

The study is reported in Nature Ecology and Evolution.

**More information:** Seasonal variation in UVA light drives hormonal and behavioral changes in a marine annelid via a ciliary opsin, *Nature Ecology and Evolution* (2021). DOI: 10.1038/s41559-020-01356-1, www.nature.com/articles/s41559-020-01356-1

Provided by University of Vienna

Citation: Ultraviolet light helps marine animals to tell the time of year (2021, January 11) retrieved 27 June 2024 from <u>https://phys.org/news/2021-01-ultraviolet-marine-animals-year.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.