


Considering disorder and cooperative effects in photon escape rates from atomic gases

February 26 2021



Credit: CC0 Public Domain

Whilst a great deal of research has studied the rates of photons escaping from cold atomic gases, these studies have used a scalar description of light leaving some of its properties untested. In a new paper published in

EPJ B Louis Bellando, a post-doctoral researcher at LOMA, University of Bordeaux, France, and his coauthors—Aharon Gero and Eric Akkermans, Technion-Israel Institute of Technology, Israel, and Robin Kaiser, Université Côte d'Azur, France—aim to numerically investigate the roles of cooperative effects and disorder in photon escape rates from a cold atomic gas to construct a model that considers the vectorial nature of light. Thus, the study accounts for properties of light, previously neglected.

"Our study focuses on light propagation in cold atomic gases, in which atoms hardly move. On their way out of the gas, photons undergo multiple scattering by the atoms," Bellando says. "Roughly speaking, the greater the number of these scattering events—the longer it takes the photons to leave the gas, and thus the smaller their escape rates. This classical description fits the so-called radiation trapping, which occurs, for example, when light undergoes a random walk in a glass of milk."

When taking into account interference and quantum mechanical effects, two mechanisms affect these escape rates: Anderson localisation arising from interference effects in the presence of disorder, and Dicke's superradiance—cooperative effects stemming from light-mediated interactions between the atoms.

Numerically studying photon escape rates from a three-dimensional cloud of cold atoms allowed the team to consider if there were any marked differences between the behavior in the simple scalar case—giving a single value to every point in a region—and the more complex vector case that assigns magnitude and direction to every point in a given area.

One of the biggest surprises encountered by the researchers as they collected their results was how well vector field observations agreed with scalar field tests. "Surprisingly, we found no significant difference

between the scalar and vectorial models, and in both cases, the dominant mechanism was cooperativity," says Bellando. "Now we know that the scalar [model](#) constitutes an excellent approximation when considering photon escape rates from atomic gases."

Because the scalar model is much simpler than the vectorial one, the similarity between the two means that in the case of [photon](#) escape rates models can use scalar fields rather than vector fields without the risk of losing substantial information.

"Light-matter interaction is an exciting field of research, both theoretically and experimentally," Bellando concludes. "Advances in this area may have a significant impact on other emerging fields, such as quantum computing."

More information: L. Bellando et al, Roles of cooperative effects and disorder in photon localization: the case of a vector radiation field, *The European Physical Journal B* (2021). [DOI: 10.1140/epjb/s10051-021-00054-6](#)

Provided by Springer

Citation: Considering disorder and cooperative effects in photon escape rates from atomic gases (2021, February 26) retrieved 12 May 2024 from <https://phys.org/news/2021-02-disorder-cooperative-effects-photon-atomic.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.