

The GRANTECAN discovers the largest cluster of galaxies known in the early universe

February 26 2021



The studied galaxy cluster in formation, 12.5 billion light years from us. The circles indicate the new members discovered with the GTC, 4 of them are shown in detail. Credit: NASA/ESA/GOODS-N+3DHST+CANDELS Team/Daniel López/IAC

A study, led by researchers at the Instituto de Astrofísica de Canarias (IAC) and carried out with OSIRIS, an instrument on the Gran Telescopio Canarias (GTC), has found the most densely populated galaxy cluster in formation in the primitive universe. The researchers predict that this structure, which is at a distance of 12.5 billion light

years from us, will have evolved becoming a cluster similar to that of Virgo, a neighbor of the Local Group of galaxies to which the Milky Way belongs. The study is published in the specialized journal *Monthly Notices of the Royal Astronomical Society (MNRAS)*.

Clusters of [galaxies](#) are groups of galaxies which remain together because of the action of gravity. To understand the evolution of these "cities of galaxies" scientists look for structures in formation, the so-called galaxy protoclusters, in the early universe.

In 2012 an international team of astronomers made an accurate determination of the distance of the galaxy HDF850.1, known as one of the galaxies with the highest rate of star formation in the observable universe. To their surprise, the scientists also discovered that this galaxy, which is one of the most studied regions on the sky, known as the Hubble Deep Field/GOODS-North, is part of a group of around a dozen protogalaxies which had formed during the first thousand million years of cosmic history. Before its discovery only one other similar primordial group was known.

Now, thanks to a new piece of research with the OSIRIS instrument on the Gran Telescopio Canarias (GTC, or GRANTECAN), the team has shown that it is one of the most densely populated regions populated with galaxies in the primitive Universe, and have for the first time carried out a detailed study of the physical properties of this system. "Surprisingly we have discovered that all the members of the cluster studied up to now, around two dozen, are galaxies with normal star formation, and that the central galaxy appears to dominate the production of stars in this structure" explains Rosa Calvi, formerly a postdoctoral researcher at the IAC and first author of the article.

Witnesses to the infancy of the local Universe

This recent study shows that this cluster of galaxies in formation is made up of various components, or "zones" with differences in their evolution. The astronomers predict that this structure will change gradually until it becomes a [galaxy cluster](#) similar to Virgo, the central region of the supercluster of the same name in which is situated the Local Group of galaxies to which the Milky Way belongs. "We see this city in construction just as it was 12,500 million years ago, when the Universe had less than 10% of its present age, so we are seeing the childhood of a cluster of galaxies like those which are typical in the local Universe" notes Helmut Dannerbauer, an IAC researcher who is co-author of this article.

The distance measured to these studied sources agrees perfectly with the predictions based on photometric observations taken previously on GRANTECAN by Pablo Arrabal Haro, formerly a doctoral student at the IAC, supervised by José Miguel Rodríguez Espinosa, an IAC researcher and Assistant General Secretary of the International Astronomical Union (IAU), and Casiana Muñoz-Tuñón, a researcher and Deputy Director of the IAC, all of them co-authors of the present article. Arrabal developed a method for selecting galaxies with normal star formation rates, based on the photometric survey SHARDS (Survey for High-z Absorption Red and Dead Sources), a Large Program of the European Southern Observatory (ESO) carried out on the GTC. "I am very happy to see that the method developed during my doctoral thesis works so well in finding and confirming a region highly populated with galaxies in the distant Universe" states Arrabal.

The SHARDS program has been led by Pablo Pérez-González, researcher at the Centro de Astrobiología (CAB, CSIC-INTA) and also author of the paper. As Pérez-González explains, "measuring exactly how these structures are forming, especially at the beginning of the Universe, is not easy, and we need exceptional data such as those we are taking with the GTC telescope as part of the SHARDS and SHARDS

Frontier Fields projects, which allow us to determine distances to galaxies and between galaxies at the edge of the Universe with a precision never achieved before."

In addition, Stefan Geier, GTC support astronomer and co-author of the paper points out that "this highly surprising result would not have been possible without the extraordinary capacity of OSIRIS together with the large collecting area of the GRANTECAN, the largest optical and infrared telescope in the world."

The Gran Telescopio Canarias and the Observatories of the Instituto de Astrofísica de Canarias (IAC) are part of the network of Singular Scientific and Technical Infrastructures of Spain.

More information: Rosa Calvi et al, Probing the existence of a rich galaxy overdensity at $z = 5.2$, *Monthly Notices of the Royal Astronomical Society* (2021). [DOI: 10.1093/mnras/staa4037](https://doi.org/10.1093/mnras/staa4037)

Provided by Instituto de Astrofísica de Canarias

Citation: The GRANTECAN discovers the largest cluster of galaxies known in the early universe (2021, February 26) retrieved 27 April 2024 from <https://phys.org/news/2021-02-grantecan-largest-cluster-galaxies-early.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--