


# Magnetic fields drive astrophysical jet shapes

February 24 2021, by Anne M Stark



This image taken with the Hubble Space Telescope shows how a bright, clumpy jet ejected from a young star has changed over time. Credit: NASA

Outflows of matter are general features stemming from systems powered by compact objects such as black holes, active galactic nuclei, pulsar wind nebulae, accreting objects such as Young Stellar Objects (YSO) and mature stars such as our sun.

But the shape of those outflows, or astrophysical jets, vary depending on

the magnetic field around them.


In new experiments, a Lawrence Livermore National Laboratory (LLNL) scientist and international collaborators found that [outflow](#) /magnetic field misalignment is a plausible key process regulating jet formation. The research appears in *Nature Communications*.

Using a high-powered laser at the École Polytechnique, the team created fast material outflows in a strong applied magnetic field as a surrogate for potential astrophysics conditions. The team specifically looked at the impact on jet formation of a misalignment between where the jet first forms and then the magnetic field.

For small misalignments, a magnetic nozzle forms and redirects the outflow in a parallel jet. For larger misalignments, this nozzle becomes increasingly asymmetric, disrupting jet formation.

"We found that outflow/magnetic field misalignment is a plausible key process regulating jet collimation in a variety of objects from our sun's outflows to extragalactic jets," said LLNL plasma physicist Drew Higginson, a co-author of the paper. "They also could provide a possible interpretation for the observed structuring of [astrophysical jets](#)."

Astrophysical jets have varied morphologies from very high aspect ratio, collimated jets, to short ones that are either clearly fragmented or are just observed and are not able to sustain a high density over a long range.



Electron density maps of 3D plasma outflow simulations with a slightly (10 degrees) and highly (45 degrees) misaligned magnetic field. This work showed that jet formation is possible with a slightly misaligned field, but not with a large misalignment. Credit: Lawrence Livermore National Laboratory

But the mechanisms underlying these varied morphologies have been unclear. In light of observations made onto a variety of astrophysical objects, the team devised a possible scenario where the relative orientation between the outflow and the large-scale ambient [magnetic field](#) surrounding the [object](#) can play a major role orienting the

dynamics of the outflow from a collimated one to a stunted, fragmented one.

**More information:** G. Revet et al. Laboratory disruption of scaled astrophysical outflows by a misaligned magnetic field, *Nature Communications* (2021). [DOI: 10.1038/s41467-021-20917-x](https://doi.org/10.1038/s41467-021-20917-x)

Provided by Lawrence Livermore National Laboratory

Citation: Magnetic fields drive astrophysical jet shapes (2021, February 24) retrieved 24 May 2024 from <https://phys.org/news/2021-02-magnetic-fields-astrophysical-jet.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--