

Giving oxygen to the question of air quality

February 18 2021



Volatile alkanes, previously believed to have only minor susceptibility to autoxidation, are now revealed to have a much more complex relationship with oxygen, which may have significant implications for fuel combustion and air pollution. Credit: © 2021 Morgan Bennett Smith.

The simplest of organic molecules have a much more complex relationship with oxygen than previously thought. Researchers from KAUST and their international collaborators have shown that alkanes participate extensively in autoxidation reactions with oxygen molecules. The discovery, which overturns current chemical wisdom, has implications for air quality prediction and efficient fuel combustion in



engines.

Autoxidation is a <u>chemical process</u> in which oxygen molecules rapidly and sequentially add to organic molecules in a radical chain reaction. The process is critical for the timing of fuel <u>combustion</u> in engines and is a key step in the atmospheric conversion of volatile organic molecules into particulate matter.

"Conventional knowledge suggests atmospheric autoxidation requires precursor <u>molecules</u> with features such as double bonds or oxygencontaining moieties," says Zhandong Wang, now a professor at the University of Science and Technology of China, formerly a research scientist with Mani Sarathy at KAUST. Alkanes—the primary component of combustion engine fuels and an important class of urban trace gases—do not have these structural features. "Alkanes were thought to have only minor susceptibility to extensive autoxidation," Wang says.

To overturn this assumption, Sarathy, Wang and colleagues showed that alkanes do undergo extensive autoxidation under the hot high-pressure conditions of combustion. The team then set out to explore the possibility that <u>alkane</u> autoxidation also occurs under <u>atmospheric</u> <u>conditions</u>.

"In 2016, we collaborated with University of Helsinki researchers to win a KAUST Competitive Research Grant," says Wang. "That was the beginning of this work."

The team used a state-of-the-art analytical technique, called chemical ionization atmospheric pressure interface time-of-flight <u>mass</u> spectrometry, to detect products of atmospheric alkane autoxidation. "Strikingly, the yield of highly oxygenated <u>organic molecules</u> containing six or more oxygen atoms was much higher than expected," Wang says.



Under combustion conditions, the team also observed alkanes that had undergone up to five sequential O_2 additions, significantly higher than the three additions they observed previously.

"These findings enrich our understanding of autoxidation processes and will allow us to better perform predictive simulations of combustion engines and atmospheric processes that impact air quality and climate," Sarathy says.

"We are now working with the department of Health, Safety and Environment in KAUST to better understand atmospheric chemical processes using real-world measurements." he adds. "Using data acquired at the KAUST-based monitoring station, we are attempting to unravel complex atmospheric chemical processes in the western region of Saudi Arabia."

The study is published in *Communications Chemistry*.

Provided by King Abdullah University of Science and Technology

Citation: Giving oxygen to the question of air quality (2021, February 18) retrieved 2 May 2024 from <u>https://phys.org/news/2021-02-oxygen-air-quality.html</u>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.