

Physicists discover new route to active matter self-organisation

February 19 2021




Figure: (a) Microscopy image of a giant vortex formed by the bacterial active matter. (b) Instantaneous velocity field of a giant vortex. Scale bar in a, b is 250

μm . (c) Periodic switching of the rotation direction of a giant vortex, oscillating like a torsional pendulum. (d) Phase space trajectory of an oscillatory giant vortex. Credit: CUHK

An international team led by Professor Yilin Wu, Associate Professor of the Department of Physics at The Chinese University of Hong Kong (CUHK) has made a novel conceptual advance in the field of active matter science. The team discovered a new route in which the self-organization of active fluids in space and time can be controlled by a single material property called viscoelasticity. This new finding may pave the way for fabricating a new class of self-driven devices and materials, such as the ability to control the rhythmic movement of soft robots without relying on electronic circuits, and for the study of microbial physiology. It has been published in the scientific journal *Nature*.

A fast growing and interdisciplinary field, active matter science studies systems consist of units where energy is spent locally to generate mechanical work. Active matter includes all living organisms from cells to animals, biopolymers driven by molecular motors, and synthetic self-propelled materials. Self-organization (the process of producing ordered structures via interaction between individual units) principles learned from these systems may find applications in [tissue engineering](#) and in fabricating new bio-inspired devices or materials.

The study was conceived by Professor Wu and his former Ph.D. student Song Liu (currently a postdoctoral fellow at the Institute for Basic Science in Korea). They have a long term interest in understanding the physic phenomena of self-organization in biological active matter, with a focus on active fluids consisting of motile microorganisms. In a previous paper co-worked with overseas physicists published in *Nature* in 2017,

they reported a weak synchronization mechanism for biological collective oscillation, in which robust temporal order emerges from a large number of erratic but weakly coupled trajectories of individual cells in bacterial suspensions. However, the simultaneous control of spatial and temporal order is more challenging.

In the new study, the CUHK research team found clues in viscoelasticity, a common property of complex fluids which have both fluid-like and solid-like responses under deformation. While manipulating the viscoelasticity of a bacterial active fluid with DNA polymers, the team found spectacular phenomena. The bacterial active fluid first self-organizes in space into a millimeter-scale rotating vortex, then displays temporal organization as the giant vortex switches its global chirality periodically with tunable frequency, like a self-driven torsional pendulum. The team believed that these striking phenomena may possibly arise from the interplay between active forcing and viscoelastic stress relaxation. Viscoelastic relaxation occurs on a [time scale](#) corresponding to the transition from solid-like to fluid-like responses when a complex fluid is deformed.

To further understand the observed phenomena, the CUHK researchers teamed up with theoretical physicists Cristina Marchetti, Professor of the University of California, Santa Barbara and her former Ph.D. student Suraj Shankar, now a Junior Fellow of Harvard University. The two theorists developed an active matter model that couples bacterial activity, polymer elastic stress, and the fields of bacterial velocity and polarization. Analysis and computer simulations of the model reproduce all the major experimental findings, and also explain the onset of spatial and temporal order in terms of the competition between the time scales of viscoelastic relaxation and active forcing.

These new findings demonstrate experimentally for the first time that viscoelasticity of materials can be harnessed to control active matter's

self-organization. It will fuel the development of non-equilibrium physics and may pave the way for fabricating a new class of adaptive self-driven devices and materials. For instance, when coupled to actuation systems of soft robots, the millimeter-scale tunable and self-oscillating vortex may be used as a 'clock generator' which provides timing signals for programmed microfluidic pumping and for controlling the rhythmic movement of soft robots, without relying on electronic circuits. Moreover, bacteria in biofilms and animal gastrointestinal tracts often swim in viscoelastic fluids abundant in long-chain polymers. The new findings also suggest that the viscoelasticity of the environment may modify the collective motion patterns of bacteria, thereby influencing the dispersal of biofilms and the translocation of gut microbiome.

More information: Song Liu et al. Viscoelastic control of spatiotemporal order in bacterial active matter, *Nature* (2021). [DOI: 10.1038/s41586-020-03168-6](https://doi.org/10.1038/s41586-020-03168-6)

Provided by The Chinese University of Hong Kong

Citation: Physicists discover new route to active matter self-organisation (2021, February 19) retrieved 25 April 2024 from <https://phys.org/news/2021-02-physicists-route-self-organisation.html>

<p>This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.</p>
--