

# Making swimming pools safer by reducing chlorine disinfection byproducts

February 17 2021



Credit: American Chemical Society

Swimming in indoor or outdoor pools is a healthy form of exercise and recreation for many people. However, studies have linked compounds that arise from chlorine disinfection of the pools to respiratory problems, including asthma, in avid swimmers. Now, researchers reporting in ACS' *Environmental Science & Technology* have found that using a complementary form of disinfection, known as copper-silver ionization (CSI), can decrease disinfection byproducts and cell toxicity of chlorinated swimming pool water.

Disinfecting swimming pool water is necessary to inactivate harmful pathogens. Although an effective disinfectant, chlorine can react with [organic matter](#) and compounds introduced by swimmers, such as those in sweat, urine, sunscreens and cosmetics, to produce disinfection byproducts (DBPs). Epidemiological studies have linked DBPs to health

problems, including respiratory issues, bladder cancer, and pregnancy and birth complications. One way to reduce the amount of added chlorine is through the use of complementary disinfection strategies, such as CSI, which involves generating antimicrobial copper and silver ions by electrolysis. Susan Richardson and colleagues wondered whether using CSI along with reduced chlorine levels could disinfect pool water while also reducing DBP formation and toxicity.

To find out, the researchers collected [water samples](#) from two pools treated with CSI and chlorine—one outdoor and one indoor. They detected 71 DBPs, some of which were quantified for the first time in pools. In experiments with mammalian cells in the lab, the team found that the indoor pool samples were more toxic to cells than the outdoor samples, likely because outdoor DBPs can volatilize in the open air or degrade with sunlight over time. In indoor [pool water](#), the lowest levels of DBP formation and toxicity were observed when the lowest amount of chlorine was used in combination with CSI. To control for factors such as number of swimmers, temperature and pH, the researchers also conducted experiments in simulated pools with a solution added that mimics human body fluids, and they observed similar results. These data suggest that using CSI with lower amounts of chlorine could be a way to make swimming safer, the researchers say.

**More information:** "Making Swimming Pools Safer: Does Copper-Silver Ionization with Chlorine Lower the Toxicity and Disinfection Byproduct Formation?" *Environmental Science & Technology* (2021). [pubs.acs.org/doi/abs/10.1021/acs.est.0c06287](https://pubs.acs.org/doi/abs/10.1021/acs.est.0c06287)

Provided by American Chemical Society

Citation: Making swimming pools safer by reducing chlorine disinfection byproducts (2021,

February 17) retrieved 5 May 2024 from <https://phys.org/news/2021-02-pools-safer-chlorine-disinfection-byproducts.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.