


# Air pollutant reductions could enhance global warming without greenhouse gas cuts

March 10 2021



Simulations using MIROC-SPRINTARS predict annual mean distribution of changes in surface air temperature following reduction of SO<sub>2</sub> emissions from fuel sources to 10% of the current level under present (left) and doubled (right) CO<sub>2</sub> concentrations. Decreasing sulfate aerosols from human activity at high CO<sub>2</sub> concentrations results in enhanced temperature increases, especially in the mid-to-high latitudes of the Northern Hemisphere.

As countries around the world race to mitigate global warming by limiting carbon dioxide emissions, an unlikely source could be making climate goals harder to achieve without even deeper cuts in greenhouse gas production: reductions in air pollution.

New modeling experiments from Kyushu University in Japan of the long-

term effects of reductions in pollutants known as [sulfate aerosols](#) predicts further increases in surface air temperature at current and increased carbon dioxide levels because of the loss of an overall cooling effect caused by the light-scattering particles.

"Air pollution causes an estimated seven million [premature deaths](#) per year worldwide, so action is essential, especially in emerging and developing countries, which tend to be most affected," says Toshihiko Takemura, professor at Kyushu University's Research Institute for Applied Mechanics and author of the study.

"However, reductions in air pollutants must come hand in hand with reductions in [greenhouse gases](#) to avoid accelerating global warming."

To analyze how sulfate aerosols—small particles of sulfur-containing compounds often produced by burning [fossil fuels](#) or biomass—influence climate, Takemura used a combination of models known as MIROC-SPRINTARS.

MIROC is a general circulation model taking into account many key aspects of the atmosphere and oceans along with their interactions, while SPRINTARS, which is widely used by news outlets for [air pollution](#) forecasts, is capable of predicting the mixing of aerosols in the atmosphere.

Combining the two models allows for effects such as the scattering and absorption of light by aerosols and the interaction of aerosols with clouds to be included in the climate projection.

Looking at the immediate changes to the atmosphere in the case of reduced emission of SO<sub>2</sub>—a precursor of sulfate aerosols—from fuel sources, Takemura found that changes such as in light scattering and cloud formation by the sulfate aerosols lead to more energy overall

entering the atmosphere, though the increase is similar regardless of whether the atmospheric carbon dioxide concentration is the same as present levels or doubled.

However, considering changes in the climate and [surface temperatures](#) over longer time scales showed that not only does the surface air temperature increase with a reduction in sulfate aerosols but this increase is even larger when carbon dioxide levels double.

"Although the fast response is similar for both situations, long-term changes caused by more slowly responding factors related to interactions with the oceans and subsequent changes, such as in clouds and precipitation, eventually leads to a bigger temperature increase," explains Takemura.

"Thus, [global warming](#) will accelerate unless increases in greenhouse gas concentrations are suppressed as air pollution control measures decrease sulfate aerosol concentrations, further emphasizing the urgency for reducing carbon dioxide in the atmosphere," he concludes.

**More information:** Toshihiko Takemura, Return to different climate states by reducing sulphate aerosols under future CO<sub>2</sub> concentrations, *Scientific Reports* (2020). [DOI: 10.1038/s41598-020-78805-1](https://doi.org/10.1038/s41598-020-78805-1)

Provided by Kyushu University

Citation: Air pollutant reductions could enhance global warming without greenhouse gas cuts (2021, March 10) retrieved 25 May 2024 from <https://phys.org/news/2021-03-air-pollutant-reductions-global-greenhouse.html>

This document is subject to copyright. Apart from any fair dealing for the purpose of private

study or research, no part may be reproduced without the written permission. The content is provided for information purposes only.